



Inden(1,2,3-cd)pyren (CAS nr. 193-39-5)
Benz(g,h,i)perylene (CAS nr. 191-24-2)

Strukturformler:



CAS nr. 193-39-5



CAS nr. 191-24-2

VKK_{ferskvand} Dækket af værdien for benz(a)pyren (*0,0082 µg/l**)

VKK_{saltvand} Dækket af værdien for benz(a)pyren (*0,00082 µg/l**)

KVKK_{ferskvand} **0,0082 µg/l****

KVKK_{saltvand} **0,00082 µg/l****

SKK_{ferskvand} **420 µg/kg tørvægt**

SKK_{saltvand} **42 µg/kg tørvægt**

BKK Dækket af værdien for benz(a)pyren

*: Hvis man har måling af disse stoffer kan disse værdier bruges. EU direktivet henviser til kvalitetskravet for benz(a)pyren. Hvis koncentrationen af B(a)P ikke overstiger kvalitetskravet regner man med at koncentrationen af disse to stoffer heller ikke overstiger miljøskadelige niveauer.

**I direktivet bruges denne værdi kun for benz(g,h,i)perylene mens der står, at KVKK ikke anvendes for Inden(1,2,3-cd)pyren.

English Summary

For benzo(ghi)perylene and indeno(123-cd)pyrene the environmental quality standards (EQS) in Directive 2008/105/EC for the water column were 0,002 µg/l for both fresh- and saltwater, and the values covered both substances. However, in the risk assessment report (RAR) on pitch coal tar, high temperature the two substances were not “pooled”, and the values for saltwater were ten times lower than the values for freshwater. Thus the values for I(1,2,3-cd)P were 0,0027 µg/l and 0,00027 µg/l for fresh- and saltwater respectively, while for B(ghi)P the values were 0,0082 µg/l and 0,00082 µg/l. The values for I(1,2,3-cd)P were set with high assessment factors (100 and 1000) because of few data. In the new directive the two substances are covered by the value for benzo(a)pyrene.

As QSAR (ECOSAR) predictions give identical results concerning the toxicity of the two substances to organisms living in water, it seems reasonable to set common quality standards for these two substances.

In the fact-sheet on the 5-6 rings PolyAromatic Hydrocarbons (PAH) from 14th of January 2011 (attached as annex (Bilag)) it is suggested to set the MAC equal to the EQS. In the new directive the MAC for B(ghi)P is 0,0082 µg/l and 0,00082 µg/l for fresh- and saltwater, respectively, whilst the MAC for I(1,2,3-cd)P is classified as “not applicable”.

Sediment quality standards (EQS_{sediment}) were derived by using the equilibrium partitioning method.

In the new directive the biota standard for protection of human health is covered by the value for benzo(a)pyrene.

The derived quality standards are:

EQS_{freshwater}	Covered by the value for benzo(a)pyrene(0,0082 µg/l*)
EQS_{saltwater}	Covered by the value for benzo(a)pyrene (0,00082 µg/l*)
MAC_{freshwater}	0,0082 µg/l**
MAC_{saltwater}	0,00082 µg/l**
EQS_{sediment, freshwater}	420 µg/kg dw
EQS_{sediment, saltwater}	42 µg/kg dw
EQS_{biota}	Covered by the value for benzo(a)pyrene

*If measurements of these two substances are available it may be possible to use these values. If not, then it is assumed that if the concentration of B(a)P does not exceed the EQS of B(a)P then the concentrations of I(1,2,3-cd)P and B(ghi)P will also not exceed harmful levels

**In the directive these values are for B(ghi)P only.

I EU's risikovurdering af tjærestofferne (CAS nr. 65996-93-2, EU-RAR, 2009) har man for CAS nr. 193-39-5 beregnet PNEC værdier på 0,0027 µg/l og 0,00027 µg/l for henholdsvis ferskvand og saltvand og for CAS nr. 191-24-2 tilsvarende beregnet henholdsvis 0,0082 µg/l og 0,00082 µg/l. Der er ikke lavet beregninger for fødekædeeffekter i hverken EU direktivet eller i EU's risikovurderingsrapport.

I det nye direktiv om kvalitetskrav (datterdirektiv til VRD) er de to stoffer ikke længere slået sammen, og der er ikke fastsat selvstændige vandkvalitetskriterier (VKK) for de to stoffer, men de dækkes af VKK for benz(a)pyren. For benz(ghi)perylene er der fastsat et KVKK, der er lig med PNEC fastsat i RAR. Der er ligeledes ikke fastsat et biotakvalitetskrav (BKK) i udkastet til nyt direktiv, idet det dækkes af BKK for benz(a)pyren.

EU-databladet til vandrammedirektivet fra januar 2011 (vedhæftet som bilag) slår heller ikke de to stoffer sammen, og konkluderer, at der er for få data for inden(1,2,3-cd)pyren til at fastsætte kvalitetskriterier, mens det følger RARen med hensyn til benz(ghi)perylene.

Begge stoffer består af 6 ringe og er relativt kompakte i strukturen og QSAR (ECOSAR) forudsigelser forudsiger samme giftighed for organismer, der lever i vand af de to stoffer. Det synes derfor rimeligt at lave fælles kvalitetskriterier for de to stoffer, som gælder for summen af dem. RARens PNEC for inden(1,2,3-cd)pyren er fastsat med en høj usikkerhedsfaktor (100 og 1000) på laveste EC₁₀.

Der er meget lille forskel på koncentrationerne for stoffernes akutte og kroniske giftighed, da de har høj akut giftighed i forbindelse med UV belysning (fototoksicitet). Derfor anvendes vandkvalitetskriteriet (VKK) som et konservativt fastsat KVKK, som det er gjort i det nye direktiv og i EU-databladet (for benz(ghi)perylene).

Sedimentkvalitetskriterier (SKK)

Der er ingen data for effekter på sedimentlevende organismer, og derfor bruges ligevægtsfordelingsmetoden (EqP).

Ifølge EU-vejledningen beregnes SKKvådvægt som følger:

$SKK_{vådvægt} = K_{susp} * VKK * 1000 / RHO_{sed}$. Faktoren 1000 er for omregning fra m³ til liter.

Ifølge RAR er $RHO_{sed} = 1150 \text{ kg/m}^3$ og $K_{susp} = 25583$ og 58607 for henholdsvis benz(ghi)perylene og inden(1,2,3-cd)pyren. Det vælges her at anvende $K_{susp} = 25583$, da det vil være en mere forsigtig tilgang.

$SKK_{vådvægt, ferskvand} = 25583 * 0.0082 \text{ µg/l} * 1000 / 1150 \text{ kg/m}^3 = 182,4 \text{ µg/kg vådvægt}$

$SKK_{tørvægt} = SKK_{vådvægt} * \text{konverteringsfaktor}$

Konverteringsfaktor = $RHO_{sed} / F_{solid} * RHO_{solid}$, hvor F_{solid} og RHO_{solid} er henholdsvis 0,2 og 2500 kg/m³ (standardværdier fra EU vejledningen).

Konverteringsfaktor = 2,3

$SKK_{\text{tørvægt, ferskvand}} = 182 \cdot 2,3 = 420 \text{ } \mu\text{g/kg tørvægt}$
 $SKK_{\text{tørvægt, saltvand}} = 420 \text{ } \mu\text{g/kg tørvægt} : 10 = 42 \text{ } \mu\text{g/kg tørvægt}$

Biotakvalitetskriterier (BKK)

I udkastet til nyt direktiv er de to stoffer dækket af BKK for benz(a)pyren

Kvalitetskriterierne bliver således:

VKK_{ferskvand} Dækket af værdien for benz(a)pyren. EU RAR: **0,0082 $\mu\text{g/l}$** .

VKK_{saltvand} Dækket af værdien for benz(a)pyren. EU RAR: **0,00082 $\mu\text{g/l}$** .

KVKK_{ferskvand} **0,0082 $\mu\text{g/l}$** . EU RAR og nye direktiv 0,0082 $\mu\text{g/l}$ for B(ghi)P men ingen værdi for I(1,2,3-cd)P

KVKK_{saltvand} **0,00082 $\mu\text{g/l}$** . RAR og nyt direktiv 0.00082 $\mu\text{g/l}$ for B(ghi)P. men ingen værdi for I(1,2,3-cd)P

SKK_{ferskvand} **420 $\mu\text{g/kg tørvægt}$**

SKK_{saltvand} **42 $\mu\text{g/kg tørvægt}$**

BKK Dækket af værdien for benz(a)pyren.

Referencer

Direktiv 2008/105/EF. Europa-Parlamentets og Rådets direktiv 2008/105/EF af 16. december 2008 om miljøkvalitetskrav inden for vandpolitikken, om ændring og senere ophævelse af Rådets direktiv 82/176/EØF, 83/513/EØF, 84/156/EØF, 84/491/EØF og 86/280/EØF og om ændring af Europa-Parlamentets og Rådets direktiv 2000/60/EF. <http://eur-lex.europa.eu/LexUriServ/LexUriServ.do?uri=OJ:L:2008:348:0084:0097:DA:PDF>

Udkast til nyt direktiv: DIRECTIVE OF THE EUROPEAN PARLIAMENT AND OF THE COUNCIL of amending Directives 2000/60/EC and 2008/105/EC as regards priority substances in the field of water policy

EU-RAR (2009). European Union Risk Assessment Report COAL-TAR PITCH, HIGH TEMPERATURE, CAS No: 65996-93-2, EINECS No: 266-028-2.

http://esis.jrc.ec.europa.eu/doc/risk_assessment/REPORT/pitchcoaltarreport323.pdf

EU vejledning: Common Implementation Strategy for the Water Framework Directive (2000/60/EC). Guidance Document No. 27. Technical Guidance for Deriving Environmental Quality Standards. <https://circabc.europa.eu/sd/d/0cc3581b-5f65-4b6f-91c6-433a1e947838/TGD-EQS%20CIS-WFD%2027%20EC%202011.pdf>

BILAG

EU-datablad fra 14. januar 2011

5-6 rings PolyAromatic Hydrocarbons (PAH)

In the currently available EQS fact sheet addressing 5-6 rings PAH, 3 EQS values are reported for the following compounds:

- Benzo[a]pyrene
- Sum of Benzo[b]fluoranthene and Benzo[k]fluoranthene
- Sum of Benzo[g,h,i]perylene and Indeno[1,2,3-cd]pyrene

These EQS are based on direct toxicity to pelagic organisms only because no data were available for protection of top predators from secondary poisoning or for protection of human health from consumption of fishery product (except for benzo[a]pyrene where a $QS_{\text{biota, hh}}$ is mentioned but not taken into account when deriving the overall QS).

Since this fact sheet was published in 2006, a final draft European Union Risk Assessment Report was made available in the context of assessment of existing chemicals (Regulation 793/93/EEC), addressing Coal Tar Pitch High Temperature (Final EU-RAR CTPHT, E.C., 2008a). By addressing CTPHT, this report provides useful information on the 5 substances addressed in the present fact sheet which are benzo[a]pyrene, benzo[b]fluoranthene, benzo[k]fluoranthene, benzo[g,h,i]perylene and indeno[1,2,3-cd]pyrene. Moreover, an unpublished report from RIVM (Verbruggen, in prep.) was provided by the Netherlands to allow a dataset as complete as possible.


Based on these new documents, an attempt was made to review the EQS for the 5 substances. While data are still missing to derive $QS_{\text{biota, sec pois}}$, a tentative $QS_{\text{biota, hh}}$ is proposed in the present document for the sum of 4 PAHs which are considered carcinogenic (Benzo[a]pyrene, Benzo[b]fluoranthene, Benzo[k]fluoranthene and Indeno[1,2,3-cd]pyrene), on the basis of an EU maximum level in foodstuffs in fish, crustaceans and molluscs. The corresponding EQS proposed for the sum of these 4 PAHs is based on this $QS_{\text{biota, hh}}$ value and is recommended for comparison with concentrations in biota, rather than in water.


A $QS_{\text{biota, hh}}$ value is also proposed for the non carcinogenic benzo[g,h,i]perylene. However, the overall QS for benzo[g,h,i]perylene is not driven by this value but by AA- $QS_{\text{water, eco}}$ value.


Following such changes, several issues remain open:


- It may be questioned whether carcinogenic substances should be pooled together to be compared to the sum of concentrations of these substances in the media while a separate EQS is proposed for non-carcinogenic substances.
- For the 4 carcinogenic PAHs, it may be questioned whether the proposed EQS should be expressed exclusively in biota or also in corresponding value in water.
- For the non-carcinogenic PAH, it may be questioned whether the proposed EQS should be expressed in water or also in biota or sediment given the high hydrophobicity potential of the substance. Finally, it may be questioned whether the proposed EQS should be expressed in total water and/or suspended particulate matter, rather than in dissolved phase of the water compartment.


1 CHEMICAL IDENTITY

Common name	Benzo[a]pyrene
Chemical name (IUPAC)	Benzo[def]chrysene
Synonym(s)	B[a]P Benzo[def]chrysene
Chemical class (when available/relevant)	Polyaromatic hydrocarbons (PAH)
CAS number	50-32-8
EC number	200-028-5
Molecular formula	C ₂₀ H ₁₂
Molecular structure	
Molecular weight (g.mol⁻¹)	252.3

Common name	Benzo[b]fluoranthene
Chemical name (IUPAC)	Benzo[e]acephenanthrylene
Synonym(s)	-
Chemical class (when available/relevant)	Polyaromatic hydrocarbons (PAH)
CAS number	205-99-2
EC number (EINECS)	205-911-9
Molecular formula	C ₂₀ H ₁₂
Molecular structure	
Molecular weight (g.mol⁻¹)	252.3

Common name	Benzo[k]fluoranthene
Chemical name (IUPAC)	Benzo[k]fluoranthene
Synonym(s)	-
Chemical class (when available/relevant)	Polyaromatic hydrocarbons (PAH)
CAS number	207-08-9
EC number (EINECS)	205-916-6
Molecular formula	C ₂₀ H ₁₂
Molecular structure	
Molecular weight (g.mol⁻¹)	252.3

Common name	Benzo[g,h,i]perylene
Chemical name (IUPAC)	Benzo[g,h,i]perylene
Synonym(s)	-
Chemical class (when available/relevant)	Polyaromatic hydrocarbons (PAH)
CAS number	191-24-2
EC number (EINECS)	205-883-8
Molecular formula	C ₂₂ H ₁₂
Molecular structure	
Molecular weight (g.mol⁻¹)	276.3

Common name	Indeno[1,2,3-cd]pyrene
Chemical name (IUPAC)	Indeno[1,2,3-cd]pyrene
Synonym(s)	-
Chemical class (when available/relevant)	Polyaromatic hydrocarbons (PAH)
CAS number	193-39-5
EC number (EINECS)	205-893-2
Molecular formula	C ₂₂ H ₁₂
Molecular structure	
Molecular weight (g.mol⁻¹)	276.3

2 EXISTING EVALUATIONS AND REGULATORY INFORMATION

Legislation	
Annex III EQS Dir. (2008/105/EC)	No (existing priority substance including in Annex I EQS Dir.)
Existing Substances Reg. (793/93/EC)	CTPHT were investigated, addressing for the purpose of assessment the PAHs studied in the present fact sheet.
Pesticides(91/414/EEC)	No
Biocides (98/8/EC)	None of the 5 substances
PBT substances	None of the 5 substances investigated by EU-PBT Working Group separately. CTPHT were investigated as a whole and it was concluded that CTPHT is considered to be a PBT and a vPvB substance.
Substances of Very High Concern (1907/2006/EC)	CTPHT are included ¹ because they are classified “Carc., PBT and vPvB” (articles 57a, 57d and 57e) Date of inclusion: 13.01.2010 Decision number ED/68/2009
POPs (Stockholm convention)	No
Other relevant chemical regulation (veterinary products, medicament, ...)	No

¹ http://echa.europa.eu/doc/candidate_list/svhc_supdoc_pitch_publication.pdf

<p>Endocrine disrupter (ED)</p> <p>E.C., 2004²</p> <p>Groshart and Okkerman, 2000</p> <p>Petersen <i>et al.</i>, 2007</p>	<ul style="list-style-type: none"> - Benzo[a]pyrene: Cat. 1 (evidence on ED) - Investigated, not categorised - Benzo[a]pyrene: * Human health: Cat. 1 (evidence on ED) - * Wildlife: Cat. 2 (potential for ED)
--	--

² Commission staff working document on implementation of the Community Strategy for Endocrine Disrupters.

3 PROPOSED QUALITY STANDARDS (QS)

3.1 ENVIRONMENTAL QUALITY STANDARD (EQS)

3.1.1 Sum of Benzo[a]pyrene, Benzo[b]fluoranthene, Benzo[k]fluoranthene and Indeno[1,2,3-cd]pyrene

Some MAC values are proposed in the following table, however, due to the high hydrophobicity of 5-6 rings PAHs, acute toxic effects are not expected to occur.

	Value	Comments
Benzo[a]pyrene		
Proposed MAC-EQS for [freshwater] [$\mu\text{g.l}^{-1}$]	0.27	See section 7.1
Proposed MAC-EQS for [marine water] [$\mu\text{g.l}^{-1}$]	0.027	
Benzo[b]fluoranthene and Benzo[k]fluoranthene		
Proposed MAC-EQS for [freshwater] [$\mu\text{g.l}^{-1}$]	0.017	See section 7.1
Proposed MAC-EQS for [marine water] [$\mu\text{g.l}^{-1}$]	0.017	
Indeno[1,2,3-cd]pyrene		
Proposed MAC-EQS for [freshwater] [$\mu\text{g.l}^{-1}$]	<i>No sufficient data available</i>	See section 7.1
Proposed MAC-EQS for [marine water] [$\mu\text{g.l}^{-1}$]		

As regards chronic effects, $QS_{\text{biota_hh}}$ for protection of human health from consumption of fishery products is deemed the “critical QS” for derivation of an Environmental Quality Standard as a first approach for the sum of the four carcinogenic PAHs, *i.e.* benzo[a]p, benzo[b]f, benzo[k]f and indeno[1,2,3-cd]p (carcinogenic PAHs). It has to be noticed however that no QS could be derived for the protection of top predators from secondary poisoning and that $QS_{\text{biota_hh}}$ is calculated on the basis of maximum levels in foodstuffs (Regulation 1881/2006/EC) (see section 7). The proposed value is as follows:

Sum of Benzo[a]pyrene, Benzo[b]fluoranthene, Benzo[k]fluoranthene and Indeno[1,2,3-cd]pyrene	Value	Comments
Proposed AA-EQS for [biota] [$\mu\text{g.kg}^{-1}_{\text{biota ww}}$]	2 for fish 5 for crustaceans and cephalopods 10 for molluscs	Critical QS is $QS_{\text{biota_hh}}$ See section 7

The proposed values correspond to concentrations in biota and are recommended for comparison with concentrations in fish or molluscs. Member States may opt to apply EQS for water instead of those recommended above. In this case, they shall convert the values recommended above in biota to their corresponding values in water by dividing them by BCF and BMF values recommended in section 5.3 of the present fact sheet.

3.1.2 Benzo[g,h,i]perylene

As regards chronic effects, AA-QS_{freshwater_eco} and AA-QS_{marine water_eco} for protection of pelagic organisms are $8.2 \cdot 10^{-3}$ and $8.2 \cdot 10^{-4} \mu\text{g.l}^{-1}$, respectively, and are deemed the “critical QS” for derivation of an Environmental Quality Standard as a first approach. The information available was sufficient to apply assessment factors of 10 and 100 to derive the AA-QS_{freshwater, eco} and AA-QS_{marine water, eco}, respectively and these values are deemed reliable.

It has to be noticed however that no QS could be derived for the protection of top predators from secondary poisoning,

As regards acute effects, the information available was sufficient to apply assessment factors of 100 and 1000, but MAC-QS_{freshwater_eco} and MAC-QS_{marine water_eco} for protection of pelagic organisms resulted in $2 \cdot 10^{-3}$ and $2 \cdot 10^{-4} \mu\text{g.l}^{-1}$, respectively. Therefore, these QS were set equal to AA-QS_{freshwater, eco} and AA-QS_{marine water, eco}, respectively.

Benzo[g,h,i]perylene	Value	Comments
Proposed AA-EQS for [freshwater] [$\mu\text{g.l}^{-1}$]	$8.2 \cdot 10^{-3}$	Critical QS is QS _{water eco}
Proposed AA-EQS in [marine waters] [$\mu\text{g.l}^{-1}$]	$8.2 \cdot 10^{-4}$	See section 7
Proposed MAC-EQS for [freshwater] [$\mu\text{g.l}^{-1}$]	$8.2 \cdot 10^{-3}$	See section 7.1
Proposed MAC-EQS for [marine water] [$\mu\text{g.l}^{-1}$]	$8.2 \cdot 10^{-4}$	

3.2 SPECIFIC QUALITY STANDARD (QS)

3.2.1 Benzo[a]pyrene

Protection objective ³	Unit	Value	Comments
Pelagic community (freshwater) – MAC-QS	[µg.l ⁻¹]	0.27	See section 7.1
Pelagic community (marine waters) – MAC-QS	[µg.l ⁻¹]	0.027	
Pelagic community (freshwater) – AA-QS	[µg.l ⁻¹]	0.022	See section 7.1
Pelagic community (marine waters) – AA-QS	[µg.l ⁻¹]	0.022	
Benthic community (freshwater)	[µg.kg ⁻¹ _{dw}]	91.5	See section 7.1
Benthic community (marine)	[µg.kg ⁻¹ _{dw}]	91.5	
Predators (secondary poisoning)	[µg.kg ⁻¹ _{biota ww}]	<i>No data available</i>	See section 7.2
	[µg.l ⁻¹]	<i>No data available</i>	
Human health via consumption of fishery products, <i>valid for the sum of benzo[a]p, benzo[b]f, benzo[k]f and indeno[1,2,3-cd]p (carcinogenic PAHs)</i>	[µg.kg ⁻¹ _{biota ww}]	- 2 for fish - 5 for crustaceans and cephalopods - 10 for molluscs	See section 7.3
	[µg.l ⁻¹]	1.7 10 ⁻⁴ (freshwater and marine waters)	
Human health via consumption of water	[µg.l ⁻¹]	0.01	

3.2.2 Benzo[b]fluoranthene

Protection objective ³	Unit	Value	Comments
Pelagic community (freshwater) – MAC-QS	[µg.l ⁻¹]	0.017	See section 7.1
Pelagic community (marine waters) – MAC-QS	[µg.l ⁻¹]	0.017	
Pelagic community (freshwater) – AA-QS	[µg.l ⁻¹]	0.017	See section 7.1
Pelagic community (marine waters) – AA-QS	[µg.l ⁻¹]	0.017	
Benthic community (freshwater)	[µg.kg ⁻¹ _{dw}]	70.7	See section 7.1
Benthic community (marine)	[µg.kg ⁻¹ _{dw}]	70.7	
Predators (secondary poisoning)	[µg.kg ⁻¹ _{biota ww}]	<i>No data available</i>	See section 7.2
	[µg.l ⁻¹]	<i>No data available</i>	
Human health via consumption of fishery products, <i>valid for the sum of benzo[a]p, benzo[b]f, benzo[k]f and indeno[1,2,3-cd]p (carcinogenic PAHs)</i>	[µg.kg ⁻¹ _{biota ww}]	- 2 for fish - 5 for crustaceans and cephalopods - 10 for molluscs	See section 7.3
	[µg.l ⁻¹]	1.7 10 ⁻⁴ (freshwater and marine waters)	
Human health via consumption of water, <i>valid for the sum of benzo[b]f, benzo[k]f, benzo[ghi]p and indeno[1,2,3-cd]p</i>	[µg.l ⁻¹]	0.1	

3.2.3 Benzo[k]fluoranthene

Protection objective ⁴	Unit	Value	Comments
-----------------------------------	------	-------	----------

³ Please note that as recommended in the Technical Guidance for deriving EQS (E.C., 2010), "EQSs [...] are not reported for 'transitional and marine waters', but either for freshwater or marine waters". If justified by substance properties or data available, QS for the different protection objectives are given independently for transitional waters or coastal and territorial waters.

Pelagic community (freshwater) – MAC-QS	[$\mu\text{g}\cdot\text{l}^{-1}$]	0.017	See section 7.1
Pelagic community (marine waters) – MAC-QS	[$\mu\text{g}\cdot\text{l}^{-1}$]	0.017	
Pelagic community (freshwater) – AA-QS	[$\mu\text{g}\cdot\text{l}^{-1}$]	0.017	See section 7.1
Pelagic community (marine water) – AA-QS	[$\mu\text{g}\cdot\text{l}^{-1}$]	0.017	
Benthic community (freshwater)	[$\mu\text{g}\cdot\text{kg}^{-1}\text{ dw}$]	67.5	See section 7.1
Benthic community (marine)	[$\mu\text{g}\cdot\text{kg}^{-1}\text{ dw}$]	67.5	
Predators (secondary poisoning)	[$\mu\text{g}\cdot\text{kg}^{-1}\text{ biota ww}$]	<i>No data available</i>	See section 7.2
	[$\mu\text{g}\cdot\text{l}^{-1}$]	<i>No data available</i>	
Human health via consumption of fishery products, <i>valid for the sum of benzo[a]p, benzo[b]f, benzo[k]f and indeno[1,2,3-cd]p (carcinogenic PAHs)</i>	[$\mu\text{g}\cdot\text{kg}^{-1}\text{ biota ww}$]	- 2 for fish - 5 for crustaceans and cephalopods - 10 for molluscs	See section 7.3
	[$\mu\text{g}\cdot\text{l}^{-1}$]	$1.7 \cdot 10^{-4}$ (freshwater and marine waters)	
Human health via consumption of water, <i>valid for the sum of benzo[b]f, benzo[k]f, benzo[ghi]p and indeno[1,2,3-cd]p</i>	[$\mu\text{g}\cdot\text{l}^{-1}$]	0.1	

3.2.4 Benzo[g,h,i]perylene

Protection objective ⁴	Unit	Value	Comments
Pelagic community (freshwater) – MAC-QS	[$\mu\text{g}\cdot\text{l}^{-1}$]	$8.2 \cdot 10^{-3}$	See section 7.1
Pelagic community (marine waters) – MAC-QS	[$\mu\text{g}\cdot\text{l}^{-1}$]	$8.2 \cdot 10^{-4}$	
Pelagic community (freshwater) – AA-QS	[$\mu\text{g}\cdot\text{l}^{-1}$]	$8.2 \cdot 10^{-3}$	See section 7.1
Pelagic community (marine water) – AA-QS	[$\mu\text{g}\cdot\text{l}^{-1}$]	$8.2 \cdot 10^{-4}$	
Benthic community (freshwater)	[$\mu\text{g}\cdot\text{kg}^{-1}\text{ dw}$]	42	See section 7.1
Benthic community (marine)	[$\mu\text{g}\cdot\text{kg}^{-1}\text{ dw}$]	4.2	
Predators (secondary poisoning)	[$\mu\text{g}\cdot\text{kg}^{-1}\text{ biota ww}$]	<i>No data available</i>	See section 7.2
	[$\mu\text{g}\cdot\text{l}^{-1}$]	<i>No data available</i>	
Human health via consumption of fishery products	[$\mu\text{g}\cdot\text{kg}^{-1}\text{ biota ww}$]	1 826	See section 7.3
	[$\mu\text{g}\cdot\text{l}^{-1}$]	0.03	
Human health via consumption of water	[$\mu\text{g}\cdot\text{l}^{-1}$]	0.1	

⁴ Please note that as recommended in the Technical Guidance for deriving EQS (E.C., 2010), "EQSs [...] are not reported for 'transitional and marine waters', but either for freshwater or marine waters". If justified by substance properties or data available, QS for the different protection objectives are given independently for transitional waters or coastal and territorial waters.

3.2.5 Indeno[1,2,3-cd]pyrene

Protection objective ⁵	Unit	Value	Comments
Pelagic community (freshwater) – MAC-QS	[$\mu\text{g.l}^{-1}$]	<i>No derivation possible</i>	See section 7.1
Pelagic community (marine waters) – MAC-QS	[$\mu\text{g.l}^{-1}$]		
Pelagic community (freshwater) – AA-QS	[$\mu\text{g.l}^{-1}$]	<i>No derivation possible</i>	See section 7.1
Pelagic community (marine water) – AA-QS	[$\mu\text{g.l}^{-1}$]		
Benthic community (freshwater)	[$\mu\text{g.kg}^{-1}_{\text{dw}}$]	<i>No derivation possible</i>	See section 7.1
Benthic community (marine)	[$\mu\text{g.kg}^{-1}_{\text{dw}}$]		
Predators (secondary poisoning)	[$\mu\text{g.kg}^{-1}_{\text{biota ww}}$]	<i>No data available</i>	See section 7.2
	[$\mu\text{g.l}^{-1}$]	<i>No data available</i>	
Human health via consumption of fishery products, <i>valid for the sum of benzo[a]p, benzo[b]f, benzo[k]f and indeno[1,2,3-cd]p (carcinogenic PAHs)</i>	[$\mu\text{g.kg}^{-1}_{\text{biota ww}}$]	- 2 for fish - 5 for crustaceans and cephalopods - 10 for molluscs	See section 7.3
	[$\mu\text{g.l}^{-1}$]	$1.7 \cdot 10^{-4}$ (freshwater and marine waters)	
Human health via consumption of water, <i>valid for the sum of benzo[b]f, benzo[k]f, benzo[ghi]p and indeno[1,2,3-cd]p</i>	[$\mu\text{g.l}^{-1}$]	0.1	

⁵ Please note that as recommended in the Technical Guidance for deriving EQS (E.C., 2010), "EQSs [...] are not reported for 'transitional and marine waters', but either for freshwater or marine waters". If justified by substance properties or data available, QS for the different protection objectives are given independently for transitional waters or coastal and territorial waters.

4 MAJOR USES AND ENVIRONMENTAL EMISSIONS

4.1 USES AND QUANTITIES

All data hereunder are extracted from Final CTPHT EU-RAR (E.C., 2008a)

4.1.1 Production

Final CTPHT EU-RAR (E.C., 2008a) states that within the European Union, high temperature coal tar pitch including 5 and 6 rings PAHs is produced “by ten companies at eleven sites in nine countries. The total European Union production capacity in 2004 was 1,127,000 tonnes. The actual production output of coal tar pitch in that year was about 817,800 tonnes. Import from outside the EU was reported to be about 91,600 tonnes per year and export was about 355,600 tonnes per year. The total consumption of coal tar pitch in the EU from these figures is estimated to be about 554,000 tonnes per year.”

4.1.2 Uses

Coal tar pitch is mainly used as a binding agent in the production of carbon electrodes, anodes and Søderberg electrodes for instance for the aluminium industry. It is also used as a binding agent for refractories, clay pigeons, active carbon, coal briquetting, road construction and roofing. Furthermore small quantities are used for heavy duty corrosion protection (see Table 2.1 of Final CTPHT EU-RAR).

4.2 ESTIMATED ENVIRONMENTAL EMISSIONS

4.2.1 Sources of PAH emissions (E.C., 2008a)

Industrial sources

“The most important industrial emission sources include coke production, primary aluminium production and creosote and wood preservation. CTPHT is produced at coke plants as such and as a by-product of primary steel production. The main source of PAH emissions in the iron and steel industry is the coke ovens, used to make coke for the steel production. (...) The coke industries improved their PAH emissions markedly by applying modern technology. Nevertheless, old installations still have high PAH emissions, leading to local high ambient air concentrations (E.C., 2001). PAH emissions at steel production using electric arc furnaces originate from the presence of tar in the used refractory material.”

“Creosote is a distillation product of coal tar, a by-product of bituminous coal coking. Emissions of PAH take place at all stages of the wood preservation process: impregnation, storage, transport and use. In the creosote and wood preservation industry, wood is mainly impregnated under pressure in vessels, but can also be sprayed or dipped. Since 2003 creosoted wood is only to be used for certain applications by professionals when treated in vacuum/pressure installations. Creosoted wood, which is treated through spraying, brushing or dipping is banned in the European Union. Creosoted wood is completely banned for certain applications like playgrounds, garden and garden furniture according to the EU Directive 2001/90/EC. Consequently wood preservation through spraying and dipping has been phased out in the European Union. Therefore emission from this source is expected to reduce considerably. PAH emissions to air from solvent use, which includes wood impregnation, in the United Kingdom clearly decreased over the period 1990 till 2002 from 104

tonnes to 69 tonnes with no clear decrease in the period 2000 till 2002. From this information it might be concluded that there is not clear direct effect on emissions from PAH resulting from the enforcement of the EU Directive at least in the United Kingdom. Other industrial sources include petrochemical and related industries (refineries), bitumen and asphalt industries (production and use), waste incineration, power plants, rubber tyre production, cement production (combustion of fossil fuels) and motor test rigs.”

Domestic sources

“PAH-emissions from domestic sources are predominantly associated with the combustion of solid fuels as wood and coal for heating and cooking purposes. These sources contribute significantly to the total PAH emission. In Europe there is a large geographic variation in these domestic emissions due to climatic differences and to the heating systems in use. In addition to heating purposes, wood, coal or peat are also burned for the decorative effect in open fireplaces.”

Mobile sources

“Mobile sources include all modes of transport using a combustion engine. PAH emissions from these sources depend on engine type, fuel type, emission control, outdoor temperature, load of vehicle, age of the car/engine and driving habits. Diesel fuelled vehicles have higher particulate emissions and the emission control equipment is less developed than gasoline vehicles. Therefore, diesel fuelled vehicles are responsible for more PAH emissions on the road. The wear and tear of tyres is also an important source of PAH emissions. Due to the extensive use of catalytic converters and improved diesel quality, the PAH emissions from tyres could even be larger than those from the exhaust of vehicles (Edlund, 2001). Non road transport includes all PAH emissions from combustion engines used by shipping activities, railways and aircrafts.”

Agricultural sources

“Agricultural sources involve the burning of organic materials under less optimum combustion activities and therefore produce significant amounts of PAH. These activities include stubble burning, open burning of land for regeneration purposes or the open burning of brushwood, trimmings, straw etc. In some EU countries there are regulations in place regulating these emissions (E.C., 2001).”

4.2.2 PAH emissions to the different compartments (E.C., 2008a)

Emissions to water

“PAH can be emitted to surface water directly or indirectly via a STP by (industrial) point sources and via atmospheric deposition. Information on PAH emission to surface water for the EU is limited to the EPER database. Based on the emission estimates for 1998 in the Netherlands, road transport is considered to be by far the largest emission source to water, followed by emissions from agriculture and consumers. The emission from industry is relatively small. The European Pollutant Emission Register (EPER, 2004) reports PAH emission of the different point sources for 2001 (see Table below). The largest industrial emission sources to water are the pre-treatment of fibres or textiles, based on the EPER data.”

PAH emissions to water in the EU for 2001 (The European Pollutant Emission Register (EPER, 2004))

Source	Direct (kg)	Indirect via STP (kg)
Industrial processes		
- Iron and steel production	10 271	381
- Petroleum industry	558	151
- Basic organic chemicals	1519	16
- Pharmaceutical products	0	36
- Pre-treatment fibres or textiles	0	12 284
Industrial combustion	1 022	6
- Installations for the production of carbon or graphite	21	0

- Slaughterhouses, plants for the production of milk other animal or vegetable raw materials	267	77
- Industrial plants for pulp from timber or other paper or board production	6	0
Waste disposal	259	80
	6	0
Total	13 923	13 031

5 ENVIRONMENTAL BEHAVIOUR

5.1 ENVIRONMENTAL DISTRIBUTION

Benzo[a]pyrene		Master reference
Water solubility (mg.l ⁻¹)	1.54 10 ⁻³	Mackay <i>et al.</i> , 1992 <i>in E.C.</i> , 2008a
Volatilisation	Benzo[a]pyrene is not likely to volatilise from surface water.	
Vapour pressure (Pa)	7.3 10 ⁻⁷ at 25°C	Mackay <i>et al.</i> , 1992 <i>in E.C.</i> , 2008a
Henry's Law constant (Pa.m ³ .mol ⁻¹)	0.034 at 20°C	Mackay <i>et al.</i> , 1992 <i>in E.C.</i> , 2008a
Adsorption	The value of 831 764 is used for derivation of QS	
Organic carbon – water partition coefficient (K _{OC})	log K _{OC} = 5.92 (<i>calculated from K_{OW}</i>) K _{OC} = 831 764	Karickhoff <i>et al.</i> , 1979
Sediment – water partition coefficient (K _{sed-water})	20 795 (<i>calculated from K_{OC}</i>)	E.C., 2010
Bioaccumulation	BCF, BMF₁ and BMF₂ values recommended for back calculation of QS_{biota} values to water reported in the dedicated following section 5.3.	
Octanol-water partition coefficient (Log K _{ow})	6.11 (<i>estimated</i>)	Mackay <i>et al.</i> , 1992 <i>in E.C.</i> , 2008a
	6.13 (<i>experimental</i>)	US-EPA, 2008

Benzo[b]fluoranthene		Master reference
Water solubility (mg.l ⁻¹)	1.28 10 ⁻³	Mackay <i>et al.</i> , 1992 <i>in E.C.</i> , 2008a
Volatilisation	Benzo[b]fluoranthene is not likely to volatilise from surface water.	
Vapour pressure (Pa)	3.3 10 ⁻⁶ at 25°C	Mackay <i>et al.</i> , 1992 <i>in E.C.</i> , 2008a
Henry's Law constant (Pa.m ³ .mol ⁻¹)	0.051 at 20°C	Mackay <i>et al.</i> , 1992 <i>in E.C.</i> , 2008a
Adsorption	The value of 831 764 is used for derivation of QS	
Organic carbon – water partition coefficient (K _{OC})	log K _{OC} = 5.92 (<i>calculated from K_{OW}</i>) K _{OC} = 831 764	Karickhoff <i>et al.</i> , 1979
Sediment – water partition coefficient (K _{sed-water})	20 795 (<i>calculated from K_{OC}</i>)	E.C., 2010
Bioaccumulation	BCF, BMF₁ and BMF₂ values recommended for back calculation of QS_{biota} values to water reported in the	

	dedicated following section 5.3..	
Octanol-water partition coefficient (Log Kow)	6.12 (<i>estimated</i>)	Mackay <i>et al.</i> , 1992 <i>in E.C.</i> , 2008a
	5.78 (<i>experimental</i>)	US-EPA, 2008

Benzo[k]fluoranthene		Master reference
Water solubility (mg.l⁻¹)	0.93 10 ⁻³	Mackay <i>et al.</i> , 1992 <i>in E.C.</i> , 2008a
Volatilisation	Benzo[k]fluoranthene is not likely to volatilise from surface water.	
Vapour pressure (Pa)	1.3 10 ⁻⁷ at 25°C	Mackay <i>et al.</i> , 1992 <i>in E.C.</i> , 2008a
Henry's Law constant (Pa.m³.mol⁻¹)	0.043 at 20°C	Mackay <i>et al.</i> , 1992 <i>in E.C.</i> , 2008a
Adsorption	The value of 794 328 is used for derivation of QS	
Organic carbon – water partition coefficient (K_{OC})	log K _{OC} = 5.9 (<i>calculated from K_{OW}</i>) K _{OC} = 794 328	Karickhoff <i>et al.</i> , 1979
Sediment – water partition coefficient (K_{sed-water})	19 859 (<i>calculated from K_{OC}</i>)	E.C., 2010
Bioaccumulation	BCF, BMF₁ and BMF₂ values recommended for back calculation of QS_{biota} values to water reported in the dedicated following section 5.3.	
Octanol-water partition coefficient (Log Kow)	6.11 (<i>estimated</i>)	Mackay <i>et al.</i> , 1992 <i>in E.C.</i> , 2008a
	6.11 (<i>experimental</i>)	US-EPA, 2008

Benzo[g,h,i]perylene		Master reference
Water solubility (mg.l⁻¹)	0.14 10 ⁻³	Mackay <i>et al.</i> , 1992 <i>in E.C.</i> , 2008a
Volatilisation	Benzo[g,h,i]perylene is not likely to volatilise from surface water.	
Vapour pressure (Pa)	1.4 10 ⁻⁸ at 25°C	Mackay <i>et al.</i> , 1992 <i>in E.C.</i> , 2008a
Henry's Law constant (Pa.m³.mol⁻¹)	0.027 at 20°C	Mackay <i>et al.</i> , 1992 <i>in E.C.</i> , 2008a
Adsorption	The value of 1 023 293 is used for derivation of QS	
Organic carbon – water partition coefficient (K_{OC})	log K _{OC} = 6.01 (<i>calculated from K_{OW}</i>) K _{OC} = 1 023 293	Karickhoff <i>et al.</i> , 1979
Sediment – water partition coefficient (K_{sed-water})	25 583 (<i>calculated from K_{OC}</i>)	E.C., 2010
Bioaccumulation	BCF, BMF₁ and BMF₂ values recommended for back calculation of QS_{biota} values to water reported in the dedicated following section 5.3.	
Octanol-water partition coefficient (Log Kow)	6.7 (<i>estimated</i>)	Mackay <i>et al.</i> , 1992 <i>in E.C.</i> , 2008a
	6.63 (<i>experimental</i>)	US-EPA, 2008

Indeno[1,2,3-cd]pyrene		Master reference
Water solubility (mg.l ⁻¹)	1 10 ⁻⁴	Mackay <i>et al.</i> , 1992 in E.C., 2008a
Volatilisation	Indeno[1,2,3-cd]pyrene is not likely to volatilise from surface water.	
Vapour pressure (Pa)	1.7 10 ⁻⁸ at 25°C	Mackay <i>et al.</i> , 1992 in E.C., 2008a
Henry's Law constant (Pa.m ³ .mol ⁻¹)	0.046 at 25°C (<i>estimated</i>)	Mackay <i>et al.</i> , 1992 in E.C., 2008a
Adsorption	The value of 1 344 229 is used for derivation of QS	
Organic carbon – water partition coefficient (K _{OC})	log K _{OC} = 6.37 (<i>calculated from K_{OW}</i>) K _{OC} = 1 344 229	Karickhoff <i>et al.</i> , 1979
Sediment – water partition coefficient (K _{sed-water})	58 607 (<i>calculated from K_{OC}</i>)	E.C., 2010
Bioaccumulation	BCF, BMF₁ and BMF₂ values recommended for back calculation of QS_{biota} values to water reported in the dedicated following section 5.3.	
Octanol-water partition coefficient (Log K _{ow})	6.7 (<i>estimated</i>)	Mackay <i>et al.</i> , 1992 in E.C., 2008a

5.2 ABIOTIC AND BIOTIC DEGRADATIONS

All information reported hereunder are extracted from Final CTPHT EU-RAR (E.C., 2008a).

Hydrolysis	PAH are chemically stable, with no functional groups that results in hydrolysis. Under environmental conditions, therefore, hydrolysis does not contribute to the degradation of PAH (Howard <i>et al.</i> , 1991).
Photolysis	The main abiotic transformation is photochemical decomposition, which in natural water takes place only in the upper few centimetres of the aqueous phase. PAHs are photodegraded by two processes, direct photolysis by light with a wavelength < 290 nm and indirect photolysis by least one oxidizing agent (Volkering and Breure, 2003). Singlet oxygen usually plays the main role in this process and the degradation process is related to the content of oxygen dissolved (Moore and Ranamoorthy, 1984). When PAHs are absorbed on particles, the accessibility for photochemical reactions may change, depending on the nature of the particles. There are great differences in photochemical reactivity between the various PAHs.

Biodegradation	<p>The results from standard test for biodegradation in water show that PAH with up to four aromatic rings are biodegradable under aerobic conditions but that the biodegradation rate of PAH with more aromatic rings is very low (EHC, 1998). Although some evidence for anaerobic transformation of PAHs has been obtained (Coates <i>et al.</i>, 1997; Thierrin <i>et al.</i>, 1993), PAHs are usually considered to be persistent under anaerobic conditions (Neff, 1979; Volkering and Breure, 2003). Because marine sediments are often anaerobic, degradation of PAHs in this compartment is expected to be very slow. The biochemical pathway for the aerobic biodegradation of PAHs has extensively been investigated. It is understood that the initial step in the aerobic catabolism of a PAH molecule by bacteria occurs via oxidation of the PAH to a dihydrodiol by a multicomponent enzyme system. These dihydroxylated intermediates may then be processed through either an ortho cleavage type of pathway, in which ring fission occurs between the two hydroxylated carbon atoms, or a meta cleavage type of pathway, which involves cleavage of the bond adjacent to the hydroxyl groups, leading to central intermediates such as protocatechates and catechols. These compounds are further converted to tricarboxylic acid cycle intermediates (van der Meer <i>et al.</i>, 1992).</p> <p>Although the biodegradation pathway of the different PAHs is very similar their biodegradation rates differ considerably. In general the biodegradation rate decreases with increasing number of aromatic rings. For example, for degradation by bacteria from estuary half lives for B[a]P of more than 1750 days was found (Gerlach, 1981). According to Volkering and Breure (2003), two factors are considered responsible for the difference in degradation rate. First, the bacterial uptake rates of the compounds with higher molecular weight have been shown to be lower than the uptake rates of the low molecular weight PAHs. The second and most important factor is the bioavailability of PAHs, due to sorption on suspended organic matter and sediment. Since the Kow and the Koc are strongly correlated, high molecular weight PAHs will degrade slower than low molecular weight PAHs. This is illustrated by Durant <i>et al.</i>, 1995 who found that the half-life of PAHs in estuarine sediment was reversely related to the Kow. Biodegradation rates also are extremely dependent on the (a)biotics conditions both in the lab and in the field. Important influencing factors are (1) the substrate concentration; with low PAH concentrations leading to longer half-lives; (2) temperature, which reversely relates to the half-live and (3) the presence or absence of a lag-phase (De Maagd, 1996). In addition, the desorption rate of PAH appears to decrease with increase of the residence time of PAHs due to slow sorption into micropores and organic matter, and polymerization or covalent binding to the organic fraction. The consequence of this aging process is a decreased biodegradability and a decreased toxicity (Volkering and Breure, 2003).</p>
-----------------------	--

5.3 BIOACCUMULATION AND BIOMAGNIFICATION POTENTIAL

		Master reference
Bioaccumulation	The BCF values of 57 981 (molluscs), 11 138 (crustaceans and cephalopods) and 135 (fish) for benzo[a]pyrene are used for derivation of $QS_{biota\ sec.\ pois.}$ for all 5-6 rings PAHs and $BMF_1 = BMF_2 = 1$ given the absence of biomagnification (Bleeker, 2009; E.C., 2010).	
BCF	<p>Values per taxa based on data reported in the BCF dedicated table below for benzo[a]pyrene:</p> <ul style="list-style-type: none"> - BCF plants = 910 (one value) - BCF annelids = 7 317 (one value) - BCF molluscs = 57 981 (geo. mean) - BCF crustaceans = 11 138 (geo. mean) - BCF insects = 1 080 (geo. mean) - BCF fish = 135 (geo. mean) <p>Moreover, these data demonstrate an absence of biomagnification given that higher trophic levels (fish) present lower BCF values than lower trophic levels such as molluscs or crustaceans. Therefore, trophic dilution seems more likely than biomagnification and BMF values should be set to 1 by default.</p> <p>Values reported for crustaceans for Benzo[k]fluoranthene and Benzo[g,h,i]perylene are of the same order of magnitude compared to Benzo[a]pyrene. The RIVM report considers that trophic dilution is relevant for these PAHs (Bleeker, 2009).</p>	Bleeker, 2009
BSAF anguilla	<p>Sum of 5 rings PAH (including Benzo[b]fluoranthene and Benzo[k]fluoranthene: 0.003 – 0.06</p> <p>Sum of 6 rings PAH (including Benzo[g,h,i]perylene and Indeno[1,2,3-cd]pyrene: 0.02 – 0.2</p>	van der Oost <i>et al.</i> , 1994 in E.C., 2008a

Table summarising BCF values for PAH 5-6 rings in several aquatic species (Bleeker, 2009)

Taxa	Species	Test system	Chem. Analysis	BCF (l.kg ⁻¹)	Type (c)	Reliability	Reference
Benzo[a]pyrene							
Pisces	<i>Lepomis macrochirus</i>	FT	¹⁴ C	367 – 608 ¹	Kin.	2	Jimenez <i>et al.</i> , 1987
		FT	¹⁴ C	30	Kin.	2	McCarthy and Jimenez, 1985
Mollusca	<i>Dreissena polymorpha</i>	S	³ H	41 000 – 84 000 ²	Kin.	2	Bruner <i>et al.</i> , 1994
		S	³ H	24 000 – 273 000 ³	Kin.	2	Gossiaux <i>et al.</i> , 1996
	<i>Perna viridis</i>	SR	GC	8 500 ⁴	Equi.	2	Richardson <i>et al.</i> , 2005
Crustacea	<i>Daphnia magna</i>	SR	HPLC	12 761	Equi.	2	Newsted and Giesy, 1987
		S	¹⁴ C	2 837	Equi.	2	Leversee <i>et al.</i> , 1981
	<i>Eurytemora affinis</i>		GCMS	1 750 ⁵	Equi.	2	Cailleaud <i>et al.</i> , 2009
	<i>Mysis relicta</i>	FT	³ H	8 496	Kin.	2	Evans and Landrum, 1989
	<i>Pontoporeia hoyi</i>	FT	¹⁴ C	73 000	Kin.	1	Landrum, 1988
			³ H	48 582	Kin.	2	Evans and Landrum, 1989
Insecta	<i>Chironomus riparius</i> (4 th instar larvae)	S	¹⁴ C	650	Equi.	2	Leversee <i>et al.</i> , 1982
		S	¹⁴ C	166	Equi.	2	Leversee <i>et al.</i> , 1981
	<i>Hexagenia limbata</i>	FT	³ H	2 725 – 11 167 ⁷	Kin.	2	Landrum and Poore, 1988
Oligochaeta	<i>Stylodrilus heringianus</i>	FT	³ H	7 317	Kin.	2	Frank <i>et al.</i> , 1986
Magnoliophyta	<i>Lemna gibba</i>	S	¹⁴ C	7 – 910 ⁶	Kin.	2	Duxbury <i>et al.</i> , 1997
Benzo[k]fluoranthene							
Crustacea	<i>Daphnia magna</i>	SR	HPLC	13 225	Equi.	2	Newsted and Giesy, 1987
Benzo[g,h,i]perylene							
Crustacea	<i>Daphnia magna</i>	SR	HPLC	28 288	Equi.	2	Newsted and Giesy, 1987

a) FT: flow-through system; S: static; SR: static renewal. b) ¹⁴C: radioactive carbon in the parent compound; GC: Gas chromatography; GCMS: Gas chromatography with mass spectrometry; Flu.Spec.: fluorescence spectrometry; ³H: radioactive hydrogen in the parent compound; HPLC: high pressure liquid chromatography. c) Kin.: Kinetic BCF, i.e. $k1/k2$; Equi.: BCF at (assumed) equilibrium, i.e. Corganism/Cwater. d) Reliability; 1: valid without restrictions; 2: valid with restrictions.

¹ BCFs were determined at different feeding regimes, i.e. fed both during uptake and depuration, not fed during uptake but fed during depuration.

² BCFs were determined with tested animals that differ in lipid content.

³ BCFs were determined at different exposure temperatures.

⁴ In this study BCF values are based on lipid weight, values given in this table are normalized to 5% lipid content.

⁵ BCFs are based on dry weight.

⁶ Values represent (a range of) BCF values from (a range of) different exposure concentrations.

6 AQUATIC ENVIRONMENTAL CONCENTRATIONS

6.1 ESTIMATED CONCENTRATIONS

As sufficient monitoring data are available no separate calculation of the regional PECs had been performed. Therefore, only C_{local} values are presenting hereunder.

6.1.1 Benzo[a]pyrene

Compartment		Predicted environmental concentration (PEC)	Master reference
Freshwater ($\mu\text{g.l}^{-1}$)	$C_{\text{local}} - \text{production}$	$1.7 \cdot 10^{-8} - 7.6 \cdot 10^{-3}$	E.C., 2008a
	$C_{\text{local}} - \text{primary Al production}$	$4.6 \cdot 10^{-4} - 0.46$	
Marine waters ($\mu\text{g.l}^{-1}$)	$C_{\text{local}} - \text{production}$	$6 \cdot 10^{-6}$	E.C., 2008a
	$C_{\text{local}} - \text{ferro-alloy producing ind.}$	0.001	
	$C_{\text{local}} - \text{primary Al production}$	$5.9 \cdot 10^{-4} - 0.36$	
Freshwater sediment ($\mu\text{g.kg}^{-1} \text{ dw}$)	$C_{\text{local}} - \text{production}$	$1.5 \cdot 10^{-3} - 736$	E.C., 2008a
	$C_{\text{local}} - \text{primary Al production}$	39 – 38 000	
Marine sediment ($\mu\text{g.kg}^{-1} \text{ dw}$)	$C_{\text{local}} - \text{production}$	0.27	E.C., 2008a
	$C_{\text{local}} - \text{ferro-alloy producing ind.}$	79.7	
	$C_{\text{local}} - \text{primary Al production}$	0.012 – 30 000	
Biota (freshwater)		No data available	
Biota (marine)		No data available	
Biota (marine predators)		No data available	

6.1.2 Benzo[b]fluoranthene

Compartment		Predicted environmental concentration (PEC)	Master reference
Freshwater ($\mu\text{g.l}^{-1}$)	$C_{\text{local}} - \text{production}$	$1.8 \cdot 10^{-8} - 1.2 \cdot 10^{-2}$	E.C., 2008a
	$C_{\text{local}} - \text{primary Al production}$	$9.8 \cdot 10^{-4} - 0.98$	
Marine waters ($\mu\text{g.l}^{-1}$)	$C_{\text{local}} - \text{production}$	$3 \cdot 10^{-6}$	E.C., 2008a
	$C_{\text{local}} - \text{ferro-alloy producing ind.}$	$1.9 \cdot 10^{-3}$	
	$C_{\text{local}} - \text{primary Al production}$	$2.5 \cdot 10^{-4} - 0.79$	
Freshwater sediment ($\mu\text{g.kg}^{-1} \text{ dw}$)	$C_{\text{local}} - \text{production}$	$1.5 \cdot 10^{-3} - 1 \cdot 10^4$	E.C., 2008a
	$C_{\text{local}} - \text{primary Al production}$	80 – 79 000	
Marine sediment ($\mu\text{g.kg}^{-1} \text{ dw}$)	$C_{\text{local}} - \text{production}$	0.23	E.C., 2008a
	$C_{\text{local}} - \text{ferro-alloy producing ind.}$	153.5	
	$C_{\text{local}} - \text{primary Al production}$	0.012 – 64 000	
Biota (freshwater)		No data available	
Biota (marine)		No data available	
Biota (marine predators)		No data available	

6.1.3 Benzo[k]fluoranthene

Compartment		Predicted environmental concentration (PEC)	Master reference
Freshwater ($\mu\text{g.l}^{-1}$)	$C_{\text{local}} - \text{production}$	$1.8 \cdot 10^{-8} - 3.8 \cdot 10^{-3}$	E.C., 2008a
Marine waters ($\mu\text{g.l}^{-1}$)	$C_{\text{local}} - \text{production}$	$3 \cdot 10^{-6}$	E.C., 2008a
Freshwater sediment ($\mu\text{g.kg}^{-1} \text{ dw}$)	$C_{\text{local}} - \text{production}$	$1.5 \cdot 10^{-3} - 345$	E.C., 2008a
Marine sediment ($\mu\text{g.kg}^{-1} \text{ dw}$)	$C_{\text{local}} - \text{production}$	0.23	E.C., 2008a
Biota (freshwater)		No data available	
Biota (marine)		No data available	
Biota (marine predators)		No data available	

6.1.4 Benzo[g,h,i]perylene

Compartment		Predicted environmental concentration (PEC)	Master reference
Freshwater ($\mu\text{g.l}^{-1}$)	$C_{\text{local}} - \text{production}$	$1.5 \cdot 10^{-8} - 4.2 \cdot 10^{-3}$	E.C., 2008a
	$C_{\text{local}} - \text{primary Al production}$	$1.6 \cdot 10^{-4} - 0.12$	
Marine waters ($\mu\text{g.l}^{-1}$)	$C_{\text{local}} - \text{production}$	$2 \cdot 10^{-6}$	E.C., 2008a
	$C_{\text{local}} - \text{ferro-alloy producing ind.}$	$4 \cdot 10^{-4}$	
	$C_{\text{local}} - \text{primary Al production}$	$1.2 \cdot 10^{-4} - 0.12$	
Freshwater sediment ($\mu\text{g.kg}^{-1} \text{ dw}$)	$C_{\text{local}} - \text{production}$	$1.6 \cdot 10^{-3} - 506$	E.C., 2008a
	$C_{\text{local}} - \text{primary Al production}$	$13 - 12\ 000$	
Marine sediment ($\mu\text{g.kg}^{-1} \text{ dw}$)	$C_{\text{local}} - \text{production}$	0.30	E.C., 2008a
	$C_{\text{local}} - \text{ferro-alloy producing ind.}$	42	
	$C_{\text{local}} - \text{primary Al production}$	$0.32 - 12\ 000$	
Biota (freshwater)		No data available	
Biota (marine)		No data available	
Biota (marine predators)		No data available	

6.1.5 Indeno[1,2,3]pyrene

Compartment		Predicted environmental concentration (PEC)	Master reference
Freshwater ($\mu\text{g.l}^{-1}$)	$C_{\text{local}} - \text{production}$	$8 \cdot 10^{-9} - 2.6 \cdot 10^{-3}$	E.C., 2008a
	$C_{\text{local}} - \text{primary Al production}$	$9.2 \cdot 10^{-4} - 0.91$	
Marine waters ($\mu\text{g.l}^{-1}$)	$C_{\text{local}} - \text{production}$	$1 \cdot 10^{-6}$	E.C., 2008a
	$C_{\text{local}} - \text{ferro-alloy producing ind.}$	$2 \cdot 10^{-4}$	
	$C_{\text{local}} - \text{primary Al production}$	$2.3 \cdot 10^{-6} - 0.071$	
Freshwater sediment ($\mu\text{g.kg}^{-1} \text{ dw}$)	$C_{\text{local}} - \text{production}$	$2 \cdot 10^{-3} - 690$	E.C., 2008a
	$C_{\text{local}} - \text{primary Al production}$	$21 - 21\ 000$	
Marine sediment ($\mu\text{g.kg}^{-1} \text{ dw}$)	$C_{\text{local}} - \text{production}$	0.37	E.C., 2008a
	$C_{\text{local}} - \text{ferro-alloy producing ind.}$	53.6	
	$C_{\text{local}} - \text{primary Al production}$	$0.55 - 17\ 000$	
Biota (freshwater)		No data available	
Biota (marine)		No data available	
Biota (marine predators)		No data available	

6.2 MEASURED CONCENTRATIONS

6.2.1 Benzo[a]pyrene

Compartment		Measured environmental concentration (MEC)	Master reference
Freshwater ($\mu\text{g.l}^{-1}$)		PEC 1: 0.035 PEC 2: 0.025	James <i>et al.</i> , 2009 ⁽¹⁾
Marine waters (coastal and/or transitional) ($\mu\text{g.l}^{-1}$)		No data available	
WWTP effluent ($\mu\text{g.l}^{-1}$)		No data available	
Sediment ($\mu\text{g.kg}^{-1} \text{ dw}$)	Sed < 2 mm	PEC 1: 300 PEC 2: 217	James <i>et al.</i> , 2009 ⁽¹⁾
	Sed 20 μm	PEC 1: 1 119 PEC 2: 1 103	
	Sed 63 μm	PEC 1: 24 PEC 2: 22	

Biota($\mu\text{g.kg}^{-1}$ ww)	Invertebrates	PEC 1: 6 PEC 2: 4	James <i>et al.</i> , 2009 ⁽¹⁾
	Fish	PEC 1: 0.014 PEC 2: 0.014	
	Marine predators	No data available	

⁽¹⁾ data originated from EU monitoring data collection

6.2.2 Benzo[b]fluoranthene

Compartment		Measured environmental concentration (MEC)	Master reference
Freshwater ($\mu\text{g.l}^{-1}$)		PEC 1: 0.036 PEC 2: 0.05	James <i>et al.</i> , 2009 ⁽¹⁾
Marine waters (coastal and/or transitional) ($\mu\text{g.l}^{-1}$)		No data available	
WWTP effluent ($\mu\text{g.l}^{-1}$)		No data available	
Sediment($\mu\text{g.kg}^{-1}$ dw)	Sed < 2 mm	PEC 1: 422 PEC 2: 310	James <i>et al.</i> , 2009 ⁽¹⁾
	Sed 20 μm	PEC 1: 1 428 PEC 2: 1 238	
	Sed 63 μm	PEC 1: 54 PEC 2: 47	
Biota($\mu\text{g.kg}^{-1}$ ww)	Invertebrates	PEC 1: 23 PEC 2: 11	James <i>et al.</i> , 2009 ⁽¹⁾
	Fish	No data available	
	Marine predators	No data available	

⁽¹⁾ data originated from EU monitoring data collection

6.2.3 Benzo[k]fluoranthene

Compartment		Measured environmental concentration (MEC)	Master reference
Freshwater ($\mu\text{g.l}^{-1}$)		PEC 1: 0.03 PEC 2: 0.025	James <i>et al.</i> , 2009 ⁽¹⁾
Marine waters (coastal and/or transitional) ($\mu\text{g.l}^{-1}$)		No data available	
WWTP effluent ($\mu\text{g.l}^{-1}$)		No data available	
Sediment($\mu\text{g.kg}^{-1}$ dw)	Sed < 2 mm	PEC 1: 219 PEC 2: 135	James <i>et al.</i> , 2009 ⁽¹⁾
	Sed 20 μm	PEC 1: 589 PEC 2: 586	
	Sed 63 μm	PEC 1: 23 PEC 2: 19	
Biota($\mu\text{g.kg}^{-1}$ ww)	Invertebrates	PEC 1: 15 PEC 2: 13	James <i>et al.</i> , 2009 ⁽¹⁾
	Fish	No data available	
	Marine predators		No data available

⁽¹⁾ data originated from EU monitoring data collection

6.2.4 Benzo[g,h,i]perylene

Compartment		Measured environmental concentration (MEC)	Master reference
Freshwater ($\mu\text{g.l}^{-1}$)		PEC 1: 0.031 PEC 2: 0.05	James <i>et al.</i> , 2009 ⁽¹⁾
Marine waters (coastal and/or transitional) ($\mu\text{g.l}^{-1}$)		No data available	
WWTP effluent ($\mu\text{g.l}^{-1}$)		No data available	
Sediment($\mu\text{g.kg}^{-1}$ dw)	Sed < 2 mm	PEC 1: 249 PEC 2: 180	James <i>et al.</i> , 2009 ⁽¹⁾
	Sed 20 μm	PEC 1: 753 PEC 2: 715	
	Sed 63 μm	PEC 1: 20 PEC 2: 19	
Biota($\mu\text{g.kg}^{-1}$ ww)	Invertebrates	PEC 1: 8 PEC 2: 7	James <i>et al.</i> , 2009 ⁽¹⁾
	Fish	No data available	
	Marine predators		No data available

⁽¹⁾ data originated from EU monitoring data collection

6.2.5 Indeno[1,2,3-cd]pyrene

Compartment		Measured environmental concentration (MEC)	Master reference
Freshwater ($\mu\text{g.l}^{-1}$)		PEC 1: 2.6 PEC 2: 0.025	James <i>et al.</i> , 2009 ⁽¹⁾
Marine waters (coastal and/or transitional) ($\mu\text{g.l}^{-1}$)		No data available	
WWTP effluent ($\mu\text{g.l}^{-1}$)		No data available	
Sediment ($\mu\text{g.kg}^{-1}$ dw)	Sed < 2 mm	PEC 1: 218 PEC 2: 171	James <i>et al.</i> , 2009 ⁽¹⁾
	Sed 20 μm	PEC 1: 740 PEC 2: 735	
	Sed 63 μm	PEC 1: 32 PEC 2: 34	
Biota ($\mu\text{g.kg}^{-1}$ ww)	Invertebrates	PEC 1: 8 PEC 2: 7	James <i>et al.</i> , 2009 ⁽¹⁾
	Fish	No data available	
	Marine predators		No data available

⁽¹⁾ data originated from EU monitoring data collection

7 EFFECTS AND QUALITY STANDARDS

Final CTPHT EU-RAR (E.C., 2008a) states that “PAHs can be toxic via different mode of actions, such as non-polar narcosis and phototoxicity. The last is caused by the ability of PAHs to absorb ultraviolet A (UVA) radiation (320–400 nm), ultraviolet B (UVB) radiation (290–320 nm), and in some instances, visible light (400–700 nm). This toxicity may occur through two mechanisms: photosensitization, and photomodification. Photosensitization generally leads to the production of singlet oxygen, a reactive oxygen species that is highly damaging to biological material. Photomodification of PAHs, usually via oxidation, results in the formation of new compounds and can occur under environmentally relevant levels of actinic radiation (Lampi *et al.*, 2005). The phototoxic effects can be observed after a short period of exposure, which explains why for PAHs like anthracene, fluoranthene and pyrene, where phototoxicity is most evident, the acute toxicity values are even lower than the chronic toxicity values. According to Weinstein and Oris (1999) there is a growing body of evidence which suggests that phototoxic PAHs may be degrading aquatic habitats, particularly those in highly contaminated areas with shallow or clear water. For example, the photoinduced chronic effects of anthracene have been reported at those UV intensities occurring at depths of 10 to 12 m in Lake Michigan (Holst & Giesy, 1989). In addition to direct uptake of PAHs from the water column, another potential route of exposure for aquatic organisms is their accumulation from sediments (see e.g. Clemens *et al.*, 1994; Kukkonen & Landrum, 1994), followed by subsequent solar ultraviolet radiation exposures closer to the surface. Ankley *et al.* (2004) also concluded in their peer review that PAHs are present at concentrations in aquatic systems such that animals can achieve tissue concentrations sufficient to cause photoactivated toxicity. Although UV penetration can vary dramatically among PAH-contaminated sites, in their view it is likely that at least some portion of the aquatic community will be exposed to UV radiation at levels sufficient to initiate photoactivated toxicity. They do recognize that at present time, the

ability to conduct PAH photoactivated risk assessment of acceptable uncertainty is limited by comprehensive information on species exposure to PAH and UV radiation during all life stages. PAH exposure and uptake, as well as UV exposure, are likely to vary considerably among species and life stages as they migrate into and out of contaminated locations and areas of high and low UV penetration. For all but sessile species, these patterns of movements are the greatest determinant of the risk for photoactivated toxicity. Despite these uncertainties, it is thought that the phototoxic effects cannot be ignored in the present risk assessment. Therefore these effects are also considered in deriving the PNECs for aquatic species. It should be noted that the UV exposure levels of the selected studies did not exceed the UV levels under natural sun light conditions.

7.1 ACUTE AND CHRONIC AQUATIC ECOTOXICITY

Ecotoxicity data reported in the tables hereunder were extracted exclusively from the finalised version of CTPHT EU-RAR (E.C., 2008a) and an RIVM report in preparation (Verbruggen, in prep.).

In the table below, all data reported were considered valid for effects assessment purpose, i.e. could be affected a reliability index (Klimisch code) of 1 or 2, or were considered useful as supporting information for effects assessment purpose, i.e. could be affected a reliability index (Klimisch code) of 2/3. Information on reliability were retrieved from finalised version of CTPHT EU-RAR (E.C., 2008a) and an RIVM report in preparation (Verbruggen, in prep.).

Averaged measured concentrations are tagged as (mm).

7.1.1 Benzo[a]pyrene

ACUTE EFFECTS – Benzo[a]pyrene

ACUTE EFFECTS – Benzo[a]pyrene			Klimisch code	Master reference
Bacteria (mg.l ⁻¹)	Freshwater	<i>Vibrio fischeri</i> / 30mm EC ₁₀ – bioluminescence > water solubility	2 acc ^{ing} to RIVM	Loibner <i>et al.</i> , 2004
	Marine			
Algae & aquatic plants (mg.l ⁻¹)	Freshwater	No information available		
	Marine	No information available		
Invertebrates (mg.l ⁻¹)	Freshwater	<i>Daphnia magna</i> (<24h) / 48h / ⁽¹⁾ EC ₅₀ >2.7 10 ⁻³ No effects observed	2 acc ^{ing} to RIVM	Bisson <i>et al.</i> , 2000b
		<i>Daphnia pulex</i> (1.9-2.1mm) / 96h / ⁽²⁾ LC ₅₀ = 5 10 ⁻³	2/3 acc ^{ing} to RIVM	Trucco <i>et al.</i> , 1983
	Marine	No information available		
	Sediment	No information available		
Fish (mg.l ⁻¹)	Freshwater	<i>Pimephales promelas</i> (larvae) / 120h / ⁽³⁾ LC ₅₀ < 5.6 10 ⁻³	2/3 acc ^{ing} to RIVM	Oris and Giesy, 1987
	Marine	No information available		
	Sediment	No information available		

⁽¹⁾ exposure in the dark

⁽²⁾ 12:12 h photoperiod, with mixed fluorescent and natural light.

⁽³⁾ LT50 study, at the end of the 96-h test period no mortality effect was found for phenanthrene and dibenzo[a,h]anthracene and less than 20% for benzo[ghi]perylene; simulated UV-A at 95 µW/m² and UV-B at 20 µW/m²; 24 h preincubation with toxicant without light; only 1 concentration tested

CHRONIC EFFECTS – Benzo[a]pyrene		Valid according to	Master reference	
Algae & aquatic plants (mg.l ⁻¹)	Freshwater	<i>Pseudokirchneriella subcapitata</i> / 72h ⁽¹⁾ EC ₁₀ – growth = 7.8 10 ⁻⁴	2 acc ^{ing} to RIVM and EU-RAR Bisson <i>et al.</i> , 2000a	
	Marine	No information available		
Invertebrates (mg.l ⁻¹)	Freshwater	<i>Ceriodaphnia dubia</i> / 7d / ⁽²⁾ EC ₁₀ – reproduction = 5 10 ⁻⁴	2 acc ^{ing} to RIVM and EU-RAR Bisson <i>et al.</i> , 2000a	
	Marine	<i>Crassostrea gigas</i> / 48h / ⁽¹⁾ NOEC _{abnormal shells} = 1 10 ⁻³	2/3 acc ^{ing} to RIVM and EU-RAR	Lyons <i>et al.</i> , 2002 in E.C., 2008a
		<i>Crassostrea gigas</i> / 48h / ⁽²⁾ NOEC _{abnormal shells} = 5 10 ⁻⁴ EC ₁₀ – abnormal shells = 2.2 10 ⁻⁴	2/3 acc ^{ing} to RIVM and EU-RAR	
		<i>Crassostrea gigas</i> / 48h / ⁽²⁾ EC ₁₀ – larval development ≥ 1.6 10 ⁻³ (mm)	2 acc ^{ing} to RIVM	AquaSense, 2004
		<i>Strongylocentrus purpuratus</i> (eggs and sperm) / 48h NOEC _{gastrula deformities} = 5 10 ⁻⁴	2 acc ^{ing} to RIVM	Hose <i>et al.</i> , 1983
	<i>Psammechinus miliaris</i> / 48h / ⁽²⁾ EC ₁₀ – larval development ≥ 1.6 10 ⁻³ (mm)	2 acc ^{ing} to RIVM	AquaSense, 2005	
	Sediment	No information available		
Fish (mg.l ⁻¹)	Freshwater	<i>Brachydanio rerio</i> / 42d / ⁽³⁾ NOEC _{ELS} ≥ 4 10 ⁻³ (one conc. tested) No effects observed	2 acc ^{ing} to RIVM Hooftman and Evers-de Ruyter, 1992	
		<i>Brachydanio rerio</i> (larvae) / 168d / ⁽³⁾ NOEC _{malformation} ≥ 4.4 10 ⁻⁴ (one conc. tested) No effects observed	2 acc ^{ing} to RIVM Petersen and Kristensen, 1998	
		<i>Oncorhynchus mykiss</i> / 36d EC ₁₀ – ELS – abnormalities ≥ 2.9 10 ⁻³	2 acc ^{ing} to RIVM Hannah <i>et al.</i> , 1982	
	Marine	<i>Psettichtys melanostichus</i> / 6d NOEC _{hatchability} < 1 10 ⁻⁴ (one conc. tested)	2/3 acc ^{ing} to RIVM Hose <i>et al.</i> , 1982	
	Sediment	No information available		

⁽¹⁾ 6000-8000 lux on the level of the solutions

⁽²⁾ 16:8 h photoperiod.

⁽³⁾ 16:8 h photoperiod, yellow light

Acute studies for freshwater species are available for crustaceans and fish. Chronic studies for freshwater species are available for algae, crustaceans, and fish. In addition, chronic studies for marine species are available for molluscs, echinoderms, and fish.

The acute test led by Bisson *et al.* (2000) on *Daphnia magna* did not result in any toxic effects. The 48h-EC₅₀ of 2.7 µg.l⁻¹ for crustacean *Daphnia magna* can however be used as endpoint for MAC-

QS_{water, eco} derivation. Assessment factors of 10 and 100 can reasonably be applied on this data to derive MAC_{freshwater, eco} and MAC_{marine water, eco}, respectively.

Many data are available that correspond to studies where no effects were observed. The EC₁₀ of 0.22 µg.l⁻¹ for shell development of the marine mollusc *Crassostrea gigas* is used as the most critical endpoint to use for AA-QS_{water, eco} derivation. Because additional chronic toxicity data are available for two groups of typical marine species, the assessment factor deemed necessary for both freshwater and marine water is 10.

Tentative QS _{water} Assessment factor method	Relevant study for derivation of QS	AF	Tentative QS
MAC _{freshwater, eco}	<i>Daphnia magna</i> / 24h	10	0.27 µg.l ⁻¹
MAC _{marine water, eco}	EC ₅₀ > 2.7 10 ⁻³ mg.l ⁻¹	100	0.027 µg.l ⁻¹
AA-QS _{freshwater, eco}	<i>Crassostrea gigas</i> / 48h	10	0.022 µg.l ⁻¹
AA-QS _{marine water, eco}	EC _{10 - abnormal shells} = 2.2 10 ⁻⁴ mg.l ⁻¹	10	0.022 µg.l ⁻¹
AA-QS _{freshwater, sed.}	No data available for sediment-dwelling organisms. Therefore, EqP method was applied to derive the AA-QS	EqP	35.2 µg.kg ⁻¹ _{ww} 91.5 µg.kg ⁻¹ _{dw}
AA-QS _{marine water, sed.}			

7.1.2 Benzo[b]fluoranthene and Benzo[k]fluoranthene

Data usable to derive the QS for benzo[b]fluoranthene are scarce. Because the two substances are mostly reported together and have structural similarities, it is proposed to combine benzo[b]fluoranthene and benzo[k]fluoranthene ecotoxicological data to derive a common QS_{water, eco}.

ACUTE EFFECTS – Benzo[b]f. and Benzo[k]f.		Klimmisch code	Master reference
Bacteria (mg.l ⁻¹)	Freshwater	No information available	
	Marine	<i>Vibrio fischeri</i> / 30mn EC _{10 - bioluminescence} > water solubility	2 acc ^{ing} to RIVM Loibner <i>et al.</i> , 2004
Algae & aquatic plants (mg.l ⁻¹)	Freshwater	No acute data are available but that “chronic data for <i>Pseudokirchneriella subcapitata</i> indicate an EC _{10 - growth - 72h} > 1 10 ⁻³ and acute data is expected to be higher”.	
	Marine	No information available	
Invertebrates (mg.l ⁻¹)	Freshwater	Data are available for [b] and [k] isomer for <i>Daphnia magna</i> only but that no effects were observed (>1.1 10 ⁻³ mg.l ⁻¹) in the two available studies (Bisson <i>et al.</i> , 2000a; Verrhiest <i>et al.</i> , 2001) and that “due to the low solubility of benzo(k)fluoranthene of about 1 µg.l ⁻¹ (Mackay and al., 2000), acute effects are not anticipated”.	
	Marine	No information available	
	Sediment	No toxicity was observed up to highest concentrations tested on <i>Rhepoxynius abronius</i> (test with benzo[b]fluoranthene), nor on <i>Hyalella azteca</i> and <i>Chironomus riparius</i> (test with benzo[k]fluoranthene)	2 acc ^{ing} to RIVM
Fish (mg.l ⁻¹)	Freshwater		
	Marine	No information available	
	Sediment	No information available	

CHRONIC EFFECTS – Benzo[b]f. and Benzo[k]f.			Valid according to	Master reference
Algae & aquatic plants (mg.l ⁻¹)	Freshwater	<i>Pseudokirchneriella subcapitata</i> / 72h / [b] and [k] EC ₁₀ – growth > 1 10 ⁻³	2 acc ^{ing} to RIVM and EU-RAR	Bisson <i>et al.</i> , 2000a
	Marine	No information available		
Invertebrates (mg.l ⁻¹)	Freshwater	<i>Ceriodaphnia dubia</i> (<24h) / 7d / [b] and [k] EC ₁₀ – reproduction > 1.08 10 ⁻³	2 acc ^{ing} to RIVM and EU-RAR	Bisson <i>et al.</i> , 2000a
		<i>Daphnia magna</i> (<24h) / 21d / [k] EC ₁₀ – mortality, offspring intrinsic growth rate > 2.2 10 ⁻³ (mm)	2 acc ^{ing} to RIVM	AquaSense, 2004
	Marine	<i>Crassostrea gigas</i> / 48h / [k] EC ₁₀ – larval development > 2.6 (mm)	2 acc ^{ing} to RIVM	AquaSense, 2004
		<i>Psammechinus miliaris</i> / 48h / [k] EC ₁₀ – larval development > 2.6 (mm)	2 acc ^{ing} to RIVM	AquaSense, 2004
	Sediment	No information available		
Fish (mg.l ⁻¹)	Freshwater	<i>Brachydanio rerio</i> / 42d / [k] EC ₁₀ – length = 1.7 10 ⁻⁴ EC ₁₀ – weight = 3.1 10 ⁻⁴	2 acc ^{ing} to RIVM and EU-RAR	Hooftman and Evers-de Ruiter, 1992
	Marine	No information available		
	Sediment	No information available		

No acute data is available for fish for benzo[b]fluoranthene and benzo[k]fluoranthene while fish is shown to be the most sensitive taxa in the chronic dataset. Therefore, it is decided to set the MAC-QS_{water, eco} at the level of the AA-QS_{water, eco}.

As regards chronic toxicity, data are available for algae, crustaceans and fish. In addition, additional marine data are available for molluscs and specific marine taxa echinoderm. The lowest EC₁₀ value is found for *Brachydanio rerio* at 0.17 µg.l⁻¹. Because additional chronic toxicity data are available for two groups of marine species, the assessment factor deemed necessary for both freshwater and marine water is 10.

Tentative QS _{water} Assessment factor method	Relevant study for derivation of QS	AF	Tentative QS	
MAC _{freshwater, eco}	No sufficient data available	-	MAC-QS is set equal to AA-QS: 0.017 µg.l ⁻¹	
MAC _{marine water, eco}		-	MAC-QS is set equal to AA-QS: 0.017 µg.l ⁻¹	
AA-QS _{freshwater, eco}	<i>Brachydanio rerio</i> / 42d	10	0.017 µg.l ⁻¹	
AA-QS _{marine water, eco}	EC ₁₀ – length = 1.7 10 ⁻⁴ mg.l ⁻¹	10	0.017 µg.l ⁻¹	
AA-QS _{freshwater, sed.}	No toxicity was observed up to highest concentrations tested. Therefore, EqP method was applied to derive the AA-QS	EqP	B[b]fluo.	B[k]fluo.
			27.2 µg.kg ⁻¹ _{ww}	26.0 µg.kg ⁻¹ _{ww}
70.7 µg.kg ⁻¹ _{dw}			67.5 µg.kg ⁻¹ _{dw}	
27.2 µg.kg ⁻¹ _{ww}			26 µg.kg ⁻¹ _{ww}	
AA-QS _{marine water, sed.}			70.7 µg.kg ⁻¹ _{dw}	67.5 µg.kg ⁻¹ _{dw}

7.1.3 Benzo[g,h,i]perylene

ACUTE EFFECTS – Benzo[g,h,i]p.

ACUTE EFFECTS – Benzo[g,h,i]p.		Klimmisch code	Master reference
Bacteria (mg.l ⁻¹)	Freshwater	No information available	
	Marine	<i>Vibrio fischeri</i> / 30mn EC ₁₀ – bioluminescence > water solubility	2 acc ^{ing} to RIVM Loibner <i>et al.</i> , 2004
Algae & aquatic plants (mg.l ⁻¹)	Freshwater	No acute data are available but chronic data for <i>Pseudokirchneriella subcapitata</i> indicate an EC _{10-growth-72h} > 1.6 10 ⁻⁴ mg.l ⁻¹ and acute data is expected to be higher.	
	Marine	No information available	
Invertebrates (mg.l ⁻¹)	Freshwater	<i>Daphnia magna</i> / 48h EC ₅₀ – immobility > 2 10 ⁻⁴ No effects observed	2 acc ^{ing} to RIVM and EU-RAR Bisson <i>et al.</i> , 2000a
	Marine	No information available	
	Sediment	No information available	
Fish (mg.l ⁻¹)	Freshwater	<i>Pimephales promelas</i> (larvae) / 120h / (1) <20% mortality at 1.5 10 ⁻⁴	2 acc ^{ing} to RIVM and EU-RAR Oris and Giesy, 1987
	Marine	No information available	
	Sediment	No information available	

(1) absence of UV-radiation and thereafter exposed to UV-light with an intensity of 20 µW/cm² UV-B (290-336 nm), 95 µW/cm² UV-A (336-400 nm). After the incubation time of 24 hours, the medium was renewed every 12 hours and exposure in combination with UVradiation lasted for 96 hours.

CHRONIC EFFECTS – Benzo[g,h,i]p.

CHRONIC EFFECTS – Benzo[g,h,i]p.		Valid according to	Master reference
Algae & aquatic plants (mg.l ⁻¹)	Freshwater	<i>Pseudokirchneriella subcapitata</i> / 72h EC ₁₀ – growth > 1.6 10 ⁻³ No effects observed	2 acc ^{ing} to RIVM Bisson <i>et al.</i> , 2000a
	Marine	No information available	
Invertebrates (mg.l ⁻¹)	Freshwater	<i>Ceriodaphnia dubia</i> (<24h) / 7d EC ₁₀ – reproduction = 8.2 10 ⁻⁵	2 acc ^{ing} to RIVM and EU-RAR Bisson <i>et al.</i> , 2000a
	Marine	No information available	
	Sediment	No information available	
Fish (mg.l ⁻¹)	Freshwater	<i>Brachydanio rerio</i> / 42d NOEC _{ELS} ≥ 1.6 10 ⁻⁴	2 acc ^{ing} to RIVM and EU-RAR Hoofman and Evers-de Rooter, 1992
	Marine	No information available	
	Sediment	No information available	

Beside the toxicity on marine bacterium *Vibrio fischeri*, no additional data on marine species are available.

Acute and chronic toxicity data are available for crustaceans and fish. Algae are not represented in the acute dataset but the chronic data show that crustaceans represent the most sensitive taxa so the dataset is considered complete.

The acute test led by Bisson *et al.* (2000) on *Daphnia magna* did not result in any toxic effects. The higher than 48h-EC₅₀ of 0.2 µg.l⁻¹ for crustacean *Daphnia magna* can however be used as endpoint for MAC-QS_{water, eco} derivation. Assessment factors of 10 and 100 can reasonably be applied on this data to derive MAC_{freshwater, eco} and MAC_{marine water, eco}, respectively.

The lowest EC₁₀ of 0.082 µg.l⁻¹ is found for reproduction of *Ceriodaphnia dubia*. Assessment factors of 10 and 100 can be applied to this data to derive the AA-QS_{freshwater, eco} and AA-QS_{marine water, eco}, respectively.

Tentative QS _{water} Assessment factor method	Relevant study for derivation of QS	AF	Tentative QS
MAC _{freshwater, eco}	<i>Daphnia magna</i> / 48h EC ₅₀ – immobility > 2 · 10 ⁻⁴ mg.l ⁻¹	100	MAC-QS = 0.002 µg.l ⁻¹ but can not be lower than AA-QS. Therefore MAC-QS is set equal to AA-QS = 8.2 · 10 ⁻³ µg.l ⁻¹
MAC _{marine water, eco}		1000	MAC-QS = 2 · 10 ⁻⁴ µg.l ⁻¹ but can not be lower than AA-QS. Therefore MAC-QS is set equal to AA-QS = 8.2 · 10 ⁻⁴ µg.l ⁻¹
AA-QS _{freshwater, eco}	<i>Ceriodaphnia dubia</i> / 7d	10	8.2 · 10 ⁻³ µg.l ⁻¹
AA-QS _{marine water, eco}	EC ₁₀ – reproduction = 8.2 · 10 ⁻⁵	100	8.2 · 10 ⁻⁴ µg.l ⁻¹
AA-QS _{freshwater, sed.}	No data available for sediment-dwelling organisms. Therefore, EqP method was applied to derive the AA-QS	EqP	16.1 µg.kg ⁻¹ _{ww} 42 µg.kg ⁻¹ _{dw}
AA-QS _{marine water, sed.}			1.6 µg.kg ⁻¹ _{ww} 4.2 µg.kg ⁻¹ _{dw}

7.1.4 Indeno[1,2,3-cd]pyrene

ACUTE EFFECTS –Indeno[1,2,3-cd]p.

		Klimmisch code	Master reference
Bacteria (mg.l ⁻¹)	Freshwater	No information available	
	Marine	<i>Vibrio fischeri</i> / 30mn EC ₁₀ – bioluminescence > water solubility	2 acc ^{ing} to RIVM Loibner <i>et al.</i> , 2004
Algae & aquatic plants (mg.l ⁻¹)	Freshwater	No information available	
	Marine	No information available	
Invertebrates (mg.l ⁻¹)	Freshwater	No information available	
	Marine	No information available	
	Sediment	No information available	
Fish (mg.l ⁻¹)	Freshwater	No information available	
	Marine		
	Sediment		

CHRONIC EFFECTS –Indeno[1,2,3-cd]p.

		Valid according to	Master reference
Algae & aquatic plants (mg.l ⁻¹)	Freshwater	<i>Pseudokirchneriella subcapitata</i> / 72h EC ₁₀ – growth = 1.5 · 10 ⁻³	2 acc ^{ing} to RIVM and EU-RAR Bisson <i>et al.</i> , 2000a
	Marine	No information available	
Invertebrates (mg.l ⁻¹)	Freshwater	<i>Ceriodaphnia dubia</i> / 7d EC ₁₀ – reproduction = 2.7 · 10 ⁻⁴	2 acc ^{ing} to RIVM and EU-RAR Bisson <i>et al.</i> , 2000a
	Marine	No information available	
	Sediment	No information available	
Fish (mg.l ⁻¹)	Freshwater	No information available	
	Marine		
	Sediment		

Acute data set for indeno[1,2,3-cd]pyrene is not sufficient to derive MAC-QS values. The chronic base-set is as well not complete for indeno[1,2,3-cd]pyrene, with missing information on chronic toxicity to fish. Therefore, no quality standard can be derived.

Tentative QS _{water} Assessment factor method	Relevant study for derivation of QS	AF	Tentative QS
MAC _{freshwater, eco}	<i>No sufficient data available</i>	-	<i>No sufficient data available</i>
MAC _{marine water, eco}		-	
AA-QS _{freshwater, eco}	<i>No sufficient data available</i>	-	<i>No sufficient data available</i>
AA-QS _{marine water, eco}		-	
AA-QS _{freshwater, sed.}	No data available for sediment-dwelling organisms.		-
AA-QS _{marine water, sed.}			-

7.2 SECONDARY POISONING

Data on the PAH toxicity to birds are scarce (Albers and Laoughlin, 2003; Patton and Dieter, 1980) and Final CTPHT EU-RAR (E.C., 2008a) states that “from these data it is not possible to derive a NOAEL for birds for either of the PAHs”. “Also relevant toxicity data to mammals is limited. Almost all of the long term studies reported were designed to assess carcinogenic potency of PAH and are not considered appropriate for secondary poisoning assessment.

Only for B[a]pyrene reprotoxicity data are available. Most severe effect were observed after administration of 10 mg.kg⁻¹ to CD-1 mice by gavage during gestation which produced decreased gonadal weights and reduced fertility and reproductive capacity in the offspring. Higher doses (40 mg.kg⁻¹) caused almost complete sterility in both sexes of offspring (Mackenzie, 1981 #7518). As no concentrations are tested a NOAEL can not be determined and consequently no QS can be derived.

Other mammalian toxicity data for Benzo[a]pyrene from 90 days studies with mice resulted in higher NOAELs (Mackenzie and Angevine, 1981). Whilst QS could be derived from these data in the usual way, the reprotoxicity data for Benzo[a]pyrene suggest that such a value might not be adequately protective and the ecological relevance of the adverse effect on which some of the NOAELs are based, might also be questionable.

Based on the available information QS_{biota, sec. pois.} values for the individual PAHs can not be derived.

Secondary poisoning of top predators		Master reference
Mammalian oral toxicity	Mice / gavage during gestation LOEC = 10 mg.kg ⁻¹ _{food}	{Mackenzie, 1981 #7518}
Avian oral toxicity	No available information	

Tentative QS _{biota secpois}	Relevant study for derivation of QS	AF	Tentative QS
Biota	No available information		

7.3 HUMAN HEALTH

Human health via consumption of fishery products				Master reference
Mammalian oral toxicity	Benzo[a]pyrene, Benzo[b]fluoranthene, Benzo[k]fluoranthene and Indeno[1,2,3-cd]pyrene are carcinogenic (cf. below). No test results usable as such are available for assessment purpose of possible effects to human <i>via</i> the consumption of fishery products for these carcinogenic substances.			
	Benzo[g,h,i]perylene is a non carcinogenic substances (cf. below). Therefore, it is deemed acceptable to use a TDI. A RIVM report (Baars <i>et al.</i> , 2001) has used the TPHCWG method (TPHCWG, 1997) ⁽¹⁾ to calculate a TPH fraction specific RfD for non carcinogenic PAHs, which is deemed applicable to benzo[g,h,i]perylene. The TL _{hh} proposed by RIVM is 0.03 mg.kg ⁻¹ .bw.d ⁻¹			
CMR ¹²	Carc.	Muta.	Repr.	
Benzo[a]pyrene	1B	1B	1B	E.C., 2008b
	1	-	-	IARC, 2009
	B2	-	-	EPA, 1986 ¹³
Benzo[b]fluoranthene	1B	nc	nc	E.C., 2008b
	2B	-	-	IARC, 2009
	B2	-	-	EPA, 1986 ¹³
Benzo[k]fluoranthene	1B	nc	nc	E.C., 2008b
	2B	-	-	IARC, 2009
	B2	-	-	EPA, 1986 ¹³
Benzo[g,h,i]perylene	nc	nc	nc	E.C., 2008b
	3	-	-	IARC, 2009
	nc	-	-	EPA, 1986 ¹³
Indeno[1,2,3-cd]pyrene	nc	nc	nc	E.C., 2008b
	2B	-	-	IARC, 2009
	B2	-	-	EPA, 1986 ¹³

For the four carcinogenic substances (Benzo[a]pyrene, Benzo[b]fluoranthene, Benzo[k]fluoranthene and Indeno[1,2,3-cd]pyrene), there are no test results available which are usable as such for assessment purpose of possible effects to human *via* the consumption of fishery products. However, Regulation 1881/2006/EC sets maximum levels for certain contaminants in foodstuffs. In Section 6 of its Annex, Polycyclic Aromatic Hydrocarbons are addressed and maximum levels are given for foodstuffs content of benzo[a]pyrene.

Maximum levels given for “fresh” (other than smoked) aquatic resources are the following:

- Fish: 2 µg.kg⁻¹_{ww}
- Crustaceans and cephalopods: 5 µg.kg⁻¹_{ww}
- Bivalve molluscs: 10 µg.kg⁻¹_{ww}

As not other data are available up to date to assess protection of human health from consumption of fishery products, it seems relevant to make use of these maximum levels to estimate a possible QS_{biota, hh}.

¹² nc: not classified; According to E.C., 2008b – Carc./Muta/Repr. 1B: presumed to have carcinogenic/mutagenic/reprotoxic effects; According to EPA, 1986 – Carc. B2: probable carcinogen; According to IARC, 2009 – 1: Carcinogenic to humans; 2B: Possibly carcinogenic to humans; 3: Not classifiable as to its carcinogenicity to humans.

¹³ U.S. EPA. 1986. Guidelines for Carcinogen Risk Assessment. 51 FR 33992-34003

Footnote of the Regulation 1881/2006/EC Annex indicates that “Benzo(a)pyrene, for which maximum levels are listed, is used as a marker for the occurrence and effect of carcinogenic polycyclic aromatic hydrocarbons. These measures therefore provide full harmonisation on polycyclic aromatic hydrocarbons in the listed foods across the Member States.”

Given that information, it is deemed relevant to use this value for the 4 carcinogenic compounds of the 5-6 rings PAH studied in this factsheet, that are Benzo[a]pyrene, Benzo[b]fluoranthene, Benzo[k]fluoranthene and Indeno[1,2,3-cd]pyrene.

For the back calculation of $QS_{\text{biota, hh}}$ into water, the BCF values of 57 981, 11 138 and 135 are used for mollusc, crustaceans and fish, respectively. BMF_1 and BMF_2 values are equal to 1 (cf. section 5.3).

Tentative $QS_{\text{biota, hh}}$	Relevant data for derivation of QS	AF	Threshold Level (mg.kg ⁻¹ _{ww})	Tentative $QS_{\text{biota, hh}}$
Human health	Maximum levels given for foodstuffs content of benzo[a]pyrene, considered applicable to sum of carcinogenic 5-6 rings PAHs which are B[a]p, B[b]fluo., B[k]fluo. and I[1,2,3-cd]pyrene: - 0.002 mg.kg ⁻¹ _{ww} for fish - 0.005 mg.kg ⁻¹ _{ww} for crustaceans and cephalopods - 0.01 mg.kg ⁻¹ _{ww} for molluscs			- fish: 2 µg.kg ⁻¹ _{ww} corresponding to 0.015 µg.l ⁻¹ - molluscs: 10 µg.kg ⁻¹ _{ww} corresponding to 1.7 10 ⁻⁴ µg.l ⁻¹ - crustaceans and cephalopods: 5 µg.kg ⁻¹ _{ww} corresponding to 4.5 10 ⁻⁴ µg.l ⁻¹
	-	-	B[g,h,i]perylene: TL _{hh} proposed by RIVM = 0.03 mg.kg ⁻¹ _{ww}	1 826 µg.kg ⁻¹ _{ww} corresponding to 0.03 µg.l ⁻¹ (using molluscs BCF) 13.5 µg.l ⁻¹ (using fish BCF)

For corresponding values in water, the worst case values of 1.7 10⁻⁴ and 0.03 µg.l⁻¹ are retained for the carcinogenic substances and benzo[g,h,i]perylene, respectively.

Human health via consumption of drinking water

Human health via consumption of drinking water		Master reference
Existing drinking water standard(s)	0.1 µg.l ⁻¹ for the sum of concentrations of: - benzo[b]fluoranthene, - benzo[k]fluoranthene, - benzo[ghi]perylene, - indeno[1,2,3-cd]pyrene.	Directive 98/83/EC
	0.01 µg.l ⁻¹ for benzo[a]pyrene	

8 BIBLIOGRAPHY, SOURCES AND SUPPORTIVE INFORMATION

Albers and Laoughlin (2003). "cited in E.C. (2008). European Union Risk Assessment Report for Coal-Tar Pitch, High Temperature (CAS-No.: 65996-93-2, EINECS-No.: 266-028-2)(Final report, Environment). Institute for Health and Consumer Protection - European Chemicals Bureau."

AquaSense (2004). Toxicity tests with priority substances in the Water Framework Directive. AquaSense, Amsterdam, The Netherlands

- AquaSense (2005). Toxicity tests with priority substances in the Water Framework Directive. AquaSense, Amsterdam, The Netherlands, Report number 2034
- Baars A.J., Theelen R.M.C., Janssen P.J.C.M., Hesse J.M., van Apeldoorn M.E., Meijerink M.C.M., Verdam L. and Zeilmaker M.J. (2001). Re-evaluation of human-toxicological maximum permissible risk levels. RIVM report 711701 025. RIVM, Bilthoven. March 2001. <http://www.rivm.nl/bibliotheek/rapporten/711701025.pdf>.
- Bisson M., Dujardin R., Flammarion P., Garric J., Babut M., Lamy M.-H., Porcher J.-M., Thybaud É. and Vindimian É. (2000a). Complément au SEQ-Eau: méthode de détermination des seuils de qualité pour les substances génotoxiques. INERIS, Verneuil-en-Halatte. 6 avril 2000. In2.
- Bisson M., Dujardin R., Flammarion P., Garric J., Babut M., Lamy M.-H., Porcher J.-M., Thybaud É. and Vindimian É. (2000b). Complément au SEQ-Eau: méthode de détermination des seuils de qualité pour les substances génotoxiques. Institut National de l'Environnement Industriel et des Risques (INERIS), Agence de l'eau Rhin-Meuse, Verneuil-en-Halatte, France
- Bleeker E.A.J. (2009). Bioaccumulation of polycyclic aromatic hydrocarbons in aquatic organisms. RIVM report 601779 002. RIVM, Bilthoven. 2009. <http://www.rivm.nl/bibliotheek/rapporten/601779002.pdf>.
- Boese B.L., Lamberson J.O., Swartz R.C., Ozretich R. and Cole F. (1998). "Photoinduced toxicity of PAHs and alkylated PAHs to a marine infaunal amphipod (*Rhepoxynius abronius*)."
Archives of Environmental Contamination and Toxicology **34**: 235-240.
- Bruner K.A., Fisher S.W. and Landrum P.F. (1994). "The Role of the Zebra Mussel, *Dreissena polymorpha*, in Contaminant Cycling: I. The Effect of Body Size and Lipid Content on the Bioconcentration of PCBs and PAHs."
Journal of Great Lakes Research **20**(4): 725-734.
- Cailleaud K., Budzinski H., Le Menach K., Souissi S. and Forget-Leray J. (2009). "Uptake and elimination of hydrophobic organic contaminants in estuarine copepods: An experimental study."
Environmental Toxicology and Chemistry **28**: 239-246.
- Coates J.D., Woodward J., Allen J., Philp P. and Lovley D.R. (1997). "Anaerobic degradation of polycyclic aromatic hydrocarbons and alkanes in petroleum-contaminated marine harbor sediments."
Applied and Environmental Microbiology **63**(9): 3589-3593.
- De Maagd P.G.J. (1996). Polycyclic aromatic hydrocarbons: fate and effects in the aquatic environment. Academic Thesis. Utrecht, The Netherlands.
- Durant N.D., Wilson L.P. and Bouwer E.J. (1995). "Microcosm studies of subsurface PAH-degrading bacteria from a former manufactured gas plant."
Journal of Contaminant Hydrology **17**(3): 213-237.
- Duxbury C.L., Dixon D.G. and Greenberg B.M. (1997). "Effects of simulated solar radiation on the bioaccumulation of polycyclic aromatic hydrocarbons by the duckweed *Lemna gibba*."
Environmental Toxicology and Chemistry **16**(8): 1739-1748.
- E.C. (2001). Ambient Air Pollution by Polycyclic Aromatic Hydrocarbons (PAH). Position Paper. Prepared by the Working Group on Polycyclic Aromatic Hydrocarbons. July, 2001.
- E.C. (2004). Commission staff working document on implementation of the Community Strategy for Endocrine Disrupters - a range of substances suspected of interfering with the hormone systems of humans and wildlife (COM(1999) 706). SEC(2004) 1372. European Commission, Brussels
- E.C. (2008a). European Union Risk Assessment Report for Coal-Tar Pitch, High Temperature (CAS-No.: 65996-93-2, EINECS-No.: 266-028-2)(Final report, Environment). Institute for Health and Consumer Protection - European Chemicals Bureau. May, 2008.

E.C. (2008b). Regulation (EC) No 1272/2008 of the European Parliament and of the Council of 16 December 2008 on classification, labelling and packaging of substances and mixtures, amending and repealing Directives 67/548/EEC and 1999/45/EC, and amending Regulation (EC) No 1907/2006 (Text with EEA relevance). Official Journal of the European Union. **L353**: 1355.

E.C. (2010). Draft Technical Guidance Document for deriving Environmental Quality Standards (January 2010 version). Not yet published

Edlund (2001). "*cited in* E.C. (2008). European Union Risk Assessment Report for Coal-Tar Pitch, High Temperature (CAS-No.: 65996-93-2, EINECS-No.: 266-028-2)(Final report, Environment). Institute for Health and Consumer Protection - European Chemicals Bureau."

EHC (1998). "*cited in* E.C. (2008). European Union Risk Assessment Report for Coal-Tar Pitch, High Temperature (CAS-No.: 65996-93-2, EINECS-No.: 266-028-2)(Final report, Environment). Institute for Health and Consumer Protection - European Chemicals Bureau."

EPER (2004). The European Pollutant Emission Register (EPER). URL: <http://www.eea.europa.eu/data-and-maps/data/eper-the-european-pollutant-emission-register-4>.

Evans M.S. and Landrum P.F. (1989). "Toxicokinetics of DDE, Benzo(a)Pyrene, and 2, 4, 5, 2', 4', 5'-Hexachlorobiphenyl in *Pontoporeia Hoyi* and *Mysis Relicta*." Journal of Great Lakes Research **15**(4): 589-600.

Frank A.P., Landrum P.F. and Eadie B.J. (1986). "Polycyclic aromatic hydrocarbon : rates of uptake, depuration, and biotransformation by lake Michigan *Stylodrilus heringianus*." Chemosphere **15**: 317-330.

Gerlach S.A. (1981). Marine pollution. Diagnosis and Therapy, Springer Verlag.

Gossiaux D.C., Landrum P.F. and Fisher S.W. (1996). "Effect of temperature on the accumulation kinetics of PAHs and PCBs in the zebra mussel, *Dreissena polymorpha*." J Great Lakes Res **22**: 379-388.

Groshart C. and Okkerman P.C. (2000). Towards the establishment of a priority list of substances for further evaluation of their role in endocrine disruption: preparation of a candidate list of substances as a basis for priority setting. Final report (incorporating corrigenda to final report dated 21 June 2000). BKH Consulting Engineers, Delft, The Netherlands; in association with TNO Nutrition and Food Research, Zeist, The Netherlands

Hannah J.B., Hose J.E., Landolt M.L., Miller B.S., Felton S.P. and Iwaoka W.T. (1982). "Benzo(a)pyrene-induced morphologic and developmental abnormalities in rainbow trout." Archives of Environmental Contamination and Toxicology **11**: 727-734.

Hooftman R.N. and Evers-de Ruyter A. (1992). Early life stage tests with *Brachydanio rerio* and several polycyclic aromatic hydrocarbons using an intermittent flow-through system (draft OECD guideline). TNO-report IMW-R 92/210. TNO Environmental and Energy Research, TNO Institute of Environmental Sciences, Delft, the Netherlands, IMW-R 92/210. 1992.

Hose J., Hannah J., DiJulio D., Landolt M., Miller B., Iwaoka W. and Felton S. (1982). "Effects of benzo [a]pyrene on early development of flatfish." Arch. Environ. Contam. Toxicol. **11**: 167-171.

Hose J.E., Puffer H.W., Oshida P.S. and Bay S.M. (1983). "Developmental and cytogenetic abnormalities induced in the purple sea urchin by environmental levels of benzo(a)pyrene." Archives of Environmental Contamination and Toxicology **12**(3): 319-325.

Howard P.H., Boethling R.S., Jarvis W.F., Meylan W.M. and Michelenko E.M. (1991). Handbook of environmental degradation rates. Chelsea, Michigan, Lewis Publishers.

IARC (2009). "Agents reviewed by the IARC Monographs. Volumes 1-100A." (1-100a).

- James A., Bonnomet V., Morin A. and Fribourg-Blanc B. (2009). Implementation of requirements on Priority substances within the Context of the Water Framework Directive. Contract N° 07010401/2008/508122/ADA/D2. Final draft prioritisation process report on monitoring-based ranking., INERIS / IOW: 58.
- Jimenez B.D., Cirno C.P. and McCarthy J.F. (1987). "Effects of feeding and temperature on uptake, elimination and metabolism of benzo(a)pyrene in the bluegill sunfish (*Lepomis macrochirus*)."
Aquatic Toxicology **10**: 41-57.
- Karickhoff S.W., Brown D.S. and Scott T.A. (1979). "Sorption of hydrophobic pollutants on natural sediments."
Water Research **13**: 241-248.
- Landrum P.F. (1988). "Toxicokinetics of organic xenobiotics in the amphipod *Pontoporeia hoyi* : Role of physiological and environmental variables."
Aquatic Toxicology **12**: 245-271.
- Landrum P.F. and Poore R. (1988). "Toxicokinetics of selected xenobiotics in *Hexagenia limbata*."
Journal of Great Lakes Research **14**: 427-437.
- Leversee G.J., Giesy J.P., Landrum P.F., Gerould S., Bowling J.W., Fannin T.E. and Haddock J.D. (1982). "Kinetics and Biotransformation of Benzo(a)pyrene in *Chironomus riparius*."
Arch. Environ. Contam. Toxicol. **11**(1): 25-31.
- Leversee G.J., Giesy J.P., Landrum P.F., Bartell S., Gerould S., Bruno M., Spacie A., Bonling J., Haddock J. and Fannin T. (1981). Disposition of benzo(a)pyrene in aquatic system components: periphyton, chironomids, daphnia, fish. In: M. Cooke and A.J. Dennis (Eds.). Analysis and Biological Fate: Polynuclear Aromatic HydroCarbons, Columbus, OH, USA, Batelle Press. pp. 357-366.
- Loibner A.P., Szolar O.H.J., Braun R. and Hirmann D. (2004). "Toxicity testing of 16 priority polycyclic aromatic hydrocarbons using Lumistox®."
Environmental Toxicology and Chemistry **23**(3): 557-564.
- Lyons B.P., Pascoe C.K. and McFadzen I.R.B. (2002). "Phototoxicity of pyrene and benzo[a]pyrene to embryo-larval stages of the pacific oyster *Crassostrea gigas*."
Marine Environmental Research **54**(3-5): 627-631.
- Mackay D. and al. e. (2000). "*cited in* E.C. (2008). European Union Risk Assessment Report for Coal-Tar Pitch, High Temperature (CAS-No.: 65996-93-2, EINECS-No.: 266-028-2)(Final report, Environment). Institute for Health and Consumer Protection - European Chemicals Bureau."
- Mackay D., Shiu W.Y. and Ma K.C. (1992). Illustrated handbook of physical-chemical properties and environmental fate for organic chemicals. Volume II. Michigan, Lewis Publishers.
- Mackenzie and Angevine (1981). "*cited in* E.C. (2008). European Union Risk Assessment Report for Coal-Tar Pitch, High Temperature (CAS-No.: 65996-93-2, EINECS-No.: 266-028-2)(Final report, Environment). Institute for Health and Consumer Protection - European Chemicals Bureau."
- McCarthy J. and Jimenez B. (1985). "Reduction in bioavailability to bluegills of polycyclic aromatic hydrocarbons bound to dissolved humic material."
Environ. Toxicol. Chem. **4**: 511-521.
- Moore J.W. and Ranamoorthy S. (1984). Organic chemicals in natural waters, Springer Verlag.
- Neff J.M. (1979). Polycyclic aromatic hydrocarbons in the aquatic environment : sources, fates and biological effects. London, Applied Sciences Publishers.
- Newsted J.L. and Giesy J.P. (1987). "Predictive models for photoinduced acute toxicity of polycyclic aromatic hydrocarbons to *Daphnia magna*, Strauss (Cladocera, crustacea)."
Environmental Toxicology and Chemistry **6**(6): 445-461.
- Oris J. and Giesy J.J. (1987). "The photo-induced toxicity of polycyclic aromatic hydrocarbons to larvae of the fathead minnow (*Pimephales promelas*)."
Chemosphere **16**(7): 1395-1404.

- Patton and Dieter (1980). "cited in E.C. (2008). European Union Risk Assessment Report for Coal-Tar Pitch, High Temperature (CAS-No.: 65996-93-2, EINECS-No.: 266-028-2)(Final report, Environment). Institute for Health and Consumer Protection - European Chemicals Bureau."
- Petersen G. and Kristensen P. (1998). "Bioaccumulation of lipophilic substances in fish early life stages." Environ Toxicol Chem **17**: 1385-1395.
- Petersen G., Rasmussen D. and Gustavson K. (2007). Study on enhancing the Endocrine Disrupter priority list with a focus on low production volume chemicals. DHI, 53559
- Richardson B.J., Tse E.S.-C., De Luca-Abbott S.B., Martin M. and Lam P.K.S. (2005). "Uptake and depuration of PAHs and chlorinated pesticides by semi-permeable membrane devices (SPMDs) and green-lipped mussels (*Perna viridis*)." Marine Pollution Bulletin **51**: 975-993.
- Thierrin J., Davis G.B., Barber C., Patterson B.M., Pribac F., Power T.R. and Lambert M. (1993). "Natural degradation rates of BTEX compounds and naphthalene in a sulphate reducing groundwater environment." Hydrol. Sc. J. **38**: 309-322.
- TPHCWG (1997). Development of fraction-specific reference doses (RfDs) and reference concentrations (RfCs) for total petroleum hydrocarbons., Amherst (MA), USA
- Trucco R., Engelhardt F. and Stacey B. (1983). "Toxicity, accumulation and clearance of aromatic hydrocarbons in *Daphnia pulex*." Environ. Pollut (Series A) **31**: 191-202.
- US-EPA (2008). EPI Suite, v.4, EPA's office of pollution prevention toxics and Syracuse Research Corporation (SRC).
- van der Meer J.R., de Vos W.M., Harayama S. and Zehnder A.J. (1992). "Molecular mechanisms of genetic adaptation to xenobiotic compounds." Microbiology and Molecular Biology Reviews **56**(4): 677-694.
- van der Oost R., van Gastel L., Worst D., Hanraads M., Satumalay K., van Schooten F.-J., Heida H. and Vermeulen N.P.E. (1994). "Biochemical markers in feral roach (*Rutilus rutilus*) in relation to the bioaccumulation of organic trace pollutants." Chemosphere **29**(4): 801-817.
- Verbruggen E.M.J. (in prep.). Environmental risk limits for polycyclic aromatic hydrocarbons (PAHs). RIVM, Bilthoven, The Netherlands., RIVM Report 711701xxx. 118.
- Verrhiest G., Clément B. and Blake G. (2001). "Single and Combined Effects of Sediment-Associated PAHs on Three Species of Freshwater Macroinvertebrates." Ecotoxicology **10**(6): 363-372.
- Volkering F. and Breure A.M. (2003). Biodegradation and general aspects of bioavailability. In: Douben (Eds.). PAHs: An Ecotoxicological Perspective, Wiley. pp. 81- 98.