

Development of Mechanistic Models

Mechanistic Model for the Northern Belt Sea

Technical documentation on biogeochemical model



Development of Mechanistic Models

Mechanistic Model for the Northern Belt Sea

Technical documentation on biogeochemical model

Prepared for Danish EPA (Miljøstyrelsen, Fyn)
Represented by Mr Harley Bundgaard Madsen, Head of Section



Eelgrass in Kertinge Nor
Photo: Peter Bondo Christensen

Project manager	Anders Chr. Erichsen & Mads Birkeland
Quality supervisor	Ian Sehested Hansen
Project number	11822245
Approval date	9. September 2020
Classification	Open

CONTENTS

1	Executive Summary	1
2	Introduction	3
3	Modelling Concept	4
3.1	Mechanistic Modelling.....	4
3.2	Model Development	4
3.3	Modelling System.....	5
4	Model Set-up.....	6
4.1	Model Domain	6
4.2	Open Boundary Conditions	7
4.3	Forcings	7
4.4	Sources	8
4.5	Initial Conditions.....	8
5	Model Calibration and Validation	9
5.1	Model Calibration Procedure	9
5.2	Presentation of Key Model Results.....	9
5.3	Model Performance.....	12
5.3.1	Calibration/Validation at Station VEJ0005790	12
5.3.2	General Calibration/Validation for Danish Waters	16
6	Conclusion.....	21
7	References.....	22

FIGURES

Figure 4-1	Top: Northern Belt Sea model bathymetry. Data originates from Kystdirektoratets 50 m bathymetry updated with satellite-derived data at shallow waters (DHI 2019e). The model includes three open boundaries connecting the model area to surrounding water bodies. A) Little Belt boundary, B) Kattegat boundary, C) Great Belt boundary.	6
Figure 4-2	Resolution of the horizontal model mesh of the NBH model.	7
Figure 4-3	Run-off sources included in the Northern Belt Sea model (white dots represent sources not included in model calculations).	8
Figure 5-1	Modelled annual average bottom water concentrations of dissolved oxygen (mg/l) for 2016.	10
Figure 5-2	Modelled annual average surface water concentrations of chlorophyll-a (mg/l) for 2016.	10
Figure 5-3	Modelled yearly average surface water concentrations of total nitrogen (mg/l) for 2016.	11
Figure 5-4	Modelled yearly average surface water concentrations of total phosphorus (mg/l) for 2016.	11
Figure 5-5	Stations within the NBH-model domain (only stations with ID).....	12
Figure 5-6	Comparison of measured and modelled concentrations of dissolved oxygen (mg/l) at station VEJ0005790 in the surface and bottom (3 m) waters. Dots represent measurements, and the solid line shows modelled data for the entire period.....	13
Figure 5-7	Comparison of measured and modelled concentrations of chlorophyll-a (mg/l) at station VEJ0005790 in surface and bottom (3 m) waters. Dots represent measurements, and solid lines show modelled data for the entire period.....	13

Figure 5-8	Comparison of measured and modelled light attenuation coefficient (K_d) at station VEJ0005790 in the surface waters. Dots represent measurements, and the solid line shows modelled data for the entire period.....	14
Figure 5-9	Measured and modelled concentrations of dissolved inorganic nitrogen (DIN) at station VEJ0005790 in the surface and bottom (3 m) waters. Dots represent measurements, and solid lines show modelled data for the entire period.....	14
Figure 5-10	Measured and modelled concentrations of dissolved inorganic phosphorus (DIP) at station VEJ0005790 in the surface and bottom (3 m) waters. Dots represent measurements, and solid lines show modelled data for the entire period.....	15
Figure 5-11	Comparison of measured and modelled concentrations of total nitrogen (TN) at station VEJ0005790 in surface and bottom (3 m) waters. Scatter data represent measurements, and solid lines show modelled data for the entire period.	15
Figure 5-12	Comparison of measured and modelled concentrations of total P (TP) at station VEJ0005790 in surface and bottom (3 m) waters. Scatter data represent measurements, and solid lines show modelled data for the entire period.	16
Figure 5-15	Bar chart illustrating all model performances evaluated against three different quality measures and all parameters. The vertical line indicates the aim of 75% being 'excellent', 'very good' or 'good'.....	16

TABLES

Table 2-1	Water bodies included in the NBH model.....	3
Table 5-1	Review of model performance at station P-Bias (%) based on measured and modelled data for the validation period 2002-2016. Blue colour indicates an 'excellent' model ($ x \leq 10\%$), dark green indicates a 'very good' model ($10 < x < 20\%$), light green indicates a 'good' model ($20 < x < 40\%$), and yellow indicates a 'poor' model ($ x > 40\%$).	18
Table 5-2	Review of model performance at station Spearman Rank Correlation (no unit) based on measured and modelled data for the validation period 2002-2016. Blue colour indicates an 'excellent' model (≥ 0.90), dark green indicates a 'very good' model (0.89-0.60), light green indicates a 'good' model (0.59-0.30), and yellow indicates a 'poor' model (< 0.30).	19
Table 5-3	Review of model performance at station Cost Function (CF, no unit) based on measured and modelled data for the validation period 2002-2016. Blue colour indicates an 'excellent' model (≤ 0.40), dark green indicates a 'very good' model (0.39-1.00), light green indicates a 'good' model (1.01-2.99), and yellow indicates a 'poor' model (≥ 3.00).	20

1 Executive Summary

The model development presented in this technical note represents the biogeochemical model development for the Northern Belt Sea (NBH-model). The NBH model is part of a larger model complex comprising several mechanistic models developed by DHI and a number of statistical models developed by Aarhus University (AU), Bioscience.

The model complex is developed with the overall aim to support the Water Framework Directive (WFD) by introducing mechanistic models in as many Danish water bodies as possible and integrating with Bayesian statistical modelling, and cross-system modelling carried out by AU, Bioscience.

Here we present the overall biogeochemical model set-up covering the Northern Belt Sea, together with a quality assessment of the model performance. This specific model includes 18 Danish water bodies, some of which are also covered by other mechanistic models:

Water Body ^{*)}	Number	Water Body	Number
Kalundborg Fjord	29	Horsens Fjord, indre	128
Nærå Strand	59	Ebeltoft Vig	141
Dalby bugt	61	Stavns Fjord	142
Lillestrand	62	Knebel Vig	144
Kerteminde Fjord	84	Kalø Vig, indre	145
Kertinge Nor	85	Norsminde Fjord	146
Vejle Fjord, ydre	122	Århus Bugt, Kalø og Begtrup Vig	147
Vejle Fjord, indre	123	Århus Bugt syd, Samsø og Nordlige Bælthav	219
Horsens Fjord, ydre	127	Nordlige Lillebælt ^{**)}	224

^{*)} Water bodies defined for the River Basin Management Plans 2015-2021

The NBH biogeochemical model builds on the developed hydrodynamic model of the Northern Belt Sea and is developed to describe the biogeochemistry within the model domain with a focus on parameters relevant for WFD, including dynamics in nutrients, phytoplankton, primary production, dissolved oxygen, organic matter and benthic vegetation.

The model quality is evaluated from three model performance measures: Percent Bias (P-Bias), Spearman Rank Correlation and Cost Function (CF). According to DHI (2019b), Model Efficiency Factor (MEF) was suggested initially, but during the model development, it was concluded that MEF is not suitable to evaluate this kind of biogeochemical models, why Cost Function (CF) is introduced. The quality measure CF was used in Erichsen and Timmermann (2017) as part of an international evaluation (Hermann et al. 2017). As described in DHI (2019b), the MEF evaluates the Root Mean Square Error (RMSE) to the standard deviation (based on measurements). As model results are compared against measurements at the exact point in time in dynamic estuarine systems with strong gradients, the MEF has proven to be unsuited (due to its dependency on entirely right timing). The CF assesses the fit/misfit between measurements and observations also normalized to the standard deviation (based on measurements) why it is decided to use this measure in the overall assessment of model performance.

Concerning the performance measures, our ambition is to have 75% of all measures (Percent Bias, Spearman Rank Correlation and Cost Function) to meet 'excellent', 'very good', or 'good' for all parameters and stations (lumped).

As can be seen from the present technical note, 81% of all data sets meet the success criteria when evaluated against the three performance measures, and 76% when assessing both annual performance and summer/winter performance of all data. The average model performance, evaluated at 12 stations within the model domain, for the biogeochemical model of the Northern Belt Sea, is summarized below:

- Model performance measures for dissolved oxygen (DO) are on average 7.5% (P-Bias), 0.8 (Spearman Rank Correlation) and 0.5 (CF). The average model performance for this parameter is categorized to be 'excellent' (P-Bias) and 'very good' (Spearman Rank Correlation and CF).
- Model performance measures for chlorophyll-a (CH) are on average 47.0% (P-Bias), 0.1 (Spearman Rank Correlation) and 1.1 (CF). The average model performance for CH is categorized to be 'very good' (CF) and 'poor' (P-Bias and Spearman Rank Correlation).
- Model performance measures for light attenuation coefficient (K_d) are on average 22.7% (P-Bias), 0.1 (Spearman Rank Correlation) and 1.1 (CF). The average model performance for K_d is categorized to be 'good' (P-Bias and CF) and 'poor' (Spearman Rank Correlation).
- Model performance measures for dissolved inorganic nitrogen (DIN) are on average 32.6% (P-Bias), 0.7 (Spearman Rank Correlation) and 0.7 (CF). The average model performance for DIN is categorized to be 'very good' (Spearman Rank Correlation and CF) and 'good' (P-Bias).
- Model performance measures for dissolved inorganic phosphorus (DIP) are on average 61.7% (P-Bias), 0.3 (Spearman Rank Correlation) and 1.4 (CF). The average model performance for this parameter is categorized to be 'good' (Spearman Rank Correlation and CF) and 'poor' (P-Bias).
- Model performance measures for total nitrogen (TN) are on average 18.5% (P-Bias), 0.5 (Spearman Rank Correlation) and 0.8 (CF). The average model performance is categorized to be 'very good' (P-Bias and CF) and 'good' (Spearman Rank Correlation).
- Model performance measures for total phosphorus (TP) are on average 14.3% (P-Bias), 0.4 (Spearman Rank Correlation) and 0.8 (CF). The average model performance for TP is categorized to be 'very good' (P-Bias and CF) and 'good' (Spearman Rank Correlation).

The details behind the above performance are available in Table 5-1, Table 5-2 and Table 5-3. Time-series comparisons are available here: rbmp2021-2027.dhigroup.com (Google Chrome only).

The ambition of meeting 'excellent', 'very good', or 'good' for 75% of all parameters and stations (lumped) has been well reached. The NBH model will be applied in the Danish waterbodies of the Northern Belt Sea. In this technical note, we conclude that the NBH biogeochemical model has been developed successfully for modelling nutrient scenarios for calculation of maximum allowable inputs (MAI).

2 Introduction

The model development presented in this technical note represents the biogeochemical model development for the Northern Belt Sea and builds on top of the NBH hydrodynamic model (DHI 2019d). Documentation on the model application will be presented in the following reports. The biogeochemical model is part of the mechanistic model complex development, which includes two regional models, three local-domain models, and six estuary specific models. The model complex is developed with the overall aim to support the Water Framework Directive (WFD) by introducing mechanistic models in as many Danish water bodies as possible and integrating with Bayesian statistical modelling and cross-system modelling carried out by AU, Bioscience.

Here we present the overall biogeochemical model set-up covering the Northern Belt Sea, together with a quality assessment of the model performance. The NBH model includes the 18 Danish water bodies listed in Table 2-1 below, some of which are covered by other mechanistic models. The location of the Danish water bodies is documented in Erichsen et al. (2019).

Table 2-1 Water bodies included in the NBH model.

Water Body ^{*)}	Number	Water Body	Number
Kalundborg Fjord	29	Horsens Fjord, indre	128
Nærå Strand	59	Ebeltoft Vig	141
Dalby bugt	61	Stavns Fjord	142
Lillestrand	62	Knebel Vig	144
Kerteminde Fjord	84	Kalø Vig, indre	145
Kertinge Nor	85	Norsminde Fjord	146
Vejle Fjord, ydre	122	Århus Bugt, Kalø og Begtrup Vig	147
Vejle Fjord, indre	123	Århus Bugt syd, Samsø og Nordlige Bælthav	219
Horsens Fjord, ydre	127	Nordlige Lillebælt ^{**)}	224

^{*)} Water bodies defined for the River Basin Management Plans 2015-2021

The biogeochemical model computes the development during the modelling period in ecological parameters, including concentrations of nutrients, dissolved oxygen, and organic matter and the Secchi depth, due to, e.g. primary production. The results represent short term changes due to specific weather events, seasonal variations and interannual trends. This project will focus on summer chlorophyll-a and parameters influencing distribution and growth of eelgrass. A detailed description of the specific state variables included in the NBH biogeochemical model can be found in DHI (2019c).

According to DHI (2019b), the quality measure Model Efficiency Factor (MEF) was suggested as a quality measure initially, but during the biogeochemical model development, it was concluded that MEF is not suitable to evaluate this kind of biogeochemical models. As described in DHI (2019b), the MEF evaluates the RMSE to the standard deviation (based on measurements). As we compare model results to the measurements at the exact point in time in a number of estuary system with strong gradients and variable dynamics, the MEF has proven to be unsuited (due to its dependency on entirely right timing). For the validation of the biogeochemical models, we have included the quality measure Cost Function (CF) in replacement of MEF (Table 5-3). The CF measure was also used in Erichsen & Timmermann (2017) and describes how the difference between measured and modelled values is related to the inherent variation in field observations.

3 Modelling Concept

3.1 Mechanistic Modelling

The present technical note represents the biogeochemical part of the model complex covering the Northern Belt Sea (NBH model). The NBH model is one model out of eleven mechanistic models developed to increase the knowledge of pressures and status in Danish marine waters and to provide tools for the Danish EPA as part of the implementation of the WFD. Mechanistic models enable dynamic descriptions of ecosystems and interactions between natural forcing and anthropogenic pressures. Hence, mechanistic models can be applied for predictions of changes in specific components, like chlorophyll-a concentrations, due to changes in, e.g. anthropogenic pressures.

The NBH model is defined as a local-domain model. The mechanistic model complex development as part of the existing projects includes two regional models, three local-domain models, and six estuary specific models:

- The regional models cover both specific Danish water bodies and regional waters, such as the North Sea and a small part of the North Atlantic, which is included in the North Sea-model, and the Baltic Sea, which is covered by the IDW-model. These models provide model results for specific water bodies but, equally important, give boundaries to local-domain models and estuary-specific models.
- Local-domain models are developed to allow for resolving most small and medium-sized water bodies in the north-western Belt Sea, the south-western Belt Sea and the water bodies in and around Smålandsfarvandet.
- Estuary-specific models are developed to allow for detailed modelling of specific estuaries.

The ecological conditions in marine waters are determined by several different natural factors like water exchange, stratification, water temperature, nutrient availability, sediment characteristics, the structure of the food web, etc. On top of that, several anthropogenic factors like nutrient loadings, fishery, etc., also impact the ecosystem and potentially the ecological status.

The model developed in this specific project aims at supporting the Danish EPA's implementation of the WFD. During this project, the models are developed to represent the period 2002-2016 evaluated against NOVANA measurements. Here we use current data on solar radiation, current nutrient loadings, etc.

After the models are developed, they will be applied for scenario modelling with changed nutrient loading to assess the Maximum Allowable nutrient Inputs (MAIs).

3.2 Model Development

The model development consists of a 3D hydrodynamic model describing the physical system (water levels, current, salinity and water temperatures), and a 3D biogeochemical (ecosystem) model describing the governing biogeochemical pelagic and benthic parameters and processes like phytoplankton, dissolved oxygen, primary production, etc. The model structure is modular, meaning that a hydrodynamic model is developed independently of the biogeochemical model (for further information about the hydrodynamic model see DHI (2019a)). A more detailed description of the biogeochemical model is available in DHI (2019c), and the underlying NBH hydrodynamic model is described in DHI (2019d).

All mechanistic models have been set up and calibrated/validated for the period 2002-2016 and reported according to the performance measures P-Bias, Spearman Rank Correlation and CF (DHI 2019b). Results from the entire modelling period are furthermore presented as time series in a WEB-tool (rbmp2021-2027.dhigroup.com, Google Chrome only) with a few examples included in section 5.3. Most data used for calibration and validation originate from the national monitoring programme NOVANA (see <http://odaforalle.au.dk> for more details). For some models and some parameters, other data are included, and the specific origin of those data will be referenced when used.

3.3 Modelling System

The biogeochemical model is based on the 3D modelling software MIKE 3 HD FM (version 2017) developed by DHI together with the numerical 3D equation solver MIKE ECO Lab to describe the relevant biogeochemical processes in the modelling system. The MIKE 3 FM modelling system is based on a flexible mesh approach with horizontal mesh elements of varying size within the modelling domain. The water column is resolved by multiple layers. The modelling system has been developed for applications within oceanographic, coastal and estuarine environments.

The scientific documentation of MIKE 3 HD FM is given in DHI (2019a).

The main components and processes determining the water quality and the response in the ecosystem (e.g. changes in eelgrass biomass) are included in the biogeochemical model. They are based on external factors (meteorology and nutrient supply). The model describes the turnover of organic material and nutrients (dissolved inorganic nitrogen, phosphate and silicate), both in the pelagic (water column) and the benthic phase (seabed or sediment). The pelagic phase includes phytoplankton and nutrients, and the benthic phase covers sediment pools of nutrients and the exchange of nutrients between the sediment and water phase. Furthermore, the benthic phase of the model describes the biomass and growth of benthic vegetation at the sea bed. The mechanisms behind the biogeochemical model are described in DHI (2019c).

4 Model Set-up

The biogeochemical model for the Northern Belt Sea (NBH-model) builds on top of the hydrodynamic model (HD) and an integrated transport model (AD). The set-up and calibration/validation of the physical Northern Belt Sea model (HD and AD) are documented in technical notes (DHI 2019d).

For the present project, the model is set up for the period 2002-2016, which means that all model input data need to cover this period.




Figure 4-1 Top: Northern Belt Sea model bathymetry. Data originates from Kystdirektoratets 50 m bathymetry updated with satellite-derived data at shallow waters (DHI 2019e). The model includes three open boundaries connecting the model area to surrounding water bodies. A) Little Belt boundary, B) Kattegat boundary, C) Great Belt boundary.

4.1 Model Domain

The model domain is determined in accordance with the area of interest of the modelling study. Also, considerations of the area of influence, being the surrounding areas that affect the area of interest, and suitable open boundary locations, affect the choice of the model domain.

The model domain includes water bodies on the east coast of Jutland (from Vejle Fjord to Ebeltoft), North coast of Fyn (from Middelfart to Kerteminde Fjord), Kalundborg Fjord and around Samsø island. The model has three open boundaries connecting the model domain to the surrounding Danish waters (Figure 4-1).

The model mesh is the representation of the model domain. More specifically, the model mesh defines the model area, the location of the open boundaries, the land-water boundaries, the horizontal and vertical model resolution (discretization), and the water depths (bathymetry) of the area. The bathymetry of the NBH model is shown in Figure 4-1, whereas Figure 4-2 shows the resolution of the horizontal mesh. For the NBH model, the horizontal mesh mainly consists of triangular elements of varying sizes, but also quadrangular elements have been applied for resolving specific deep channels (i.e. the Great Belt). The horizontal resolution varies gradually

from 40-500 m in the coastal areas to 0.5-2 km in open waters. The vertical mesh is structured and consists of a combination of 10 sigma layers down to -10m and z-layers of 1m thickness for the rest of the water column. Further documentation on model mesh and horizontal/vertical resolution of the NBH HD model can be found in DHI (2019d).




Figure 4-2 Resolution of the horizontal model mesh of the NBH model.

4.2 Open Boundary Conditions

The NBH model has three open boundaries: one towards Kattegat in the east; one connecting to Great Belt in the southeast and one connecting to the Little Belt in the west (see Figure 4-1). The biogeochemical data for the boundary condition are extracted from DHI's operational Inner Danish Waters model (the DKBS2-HD75-EU model). Documentation on boundary conditions for the biogeochemical model development is given in DHI (2020).

4.3 Forcings

Data on solar radiation is calculated from clearness percentages and applied as a spatial and temporally varying forcing covering the entire NBH-model domain.

Area distributed atmospheric deposition of nitrogen (N) is provided by AU, Department of Environmental Science, and aligned with HELCOM depositions (see DHI 2020).

For estimation of suspended sediment concentrations, a dynamic bottom shear stress information is needed. Wave parameters from a Spectral Wave model are included as model forcing, including significant wave height, wave period and mean wave direction, together with current conditions from the hydrodynamic model results.

Documentation on model forcing is given in DHI (2020).

4.4 Sources

The NBH model includes sources with land-based nutrient loadings. In Figure 4-3, the location of the sources is illustrated. Freshwater run-off from land is included in the hydrodynamic module. In Denmark, 4th order area run-off distributed to main rivers and streams is applied.

The model sources are specified as time series with daily loadings of inorganic and organic nutrients, including also total nitrogen (TN) and total phosphorus (TP). The land-based nutrient loadings are based on the following data sources:

- DCE (Danish Centre for Environment and Energy) – Danish run-off

More details are included in DHI (2020).




Figure 4-3 Run-off sources included in the Northern Belt Sea model (white dots represent sources not included in model calculations).

4.5 Initial Conditions

To properly initiate a model simulation, the model requires initial conditions for the various state variables.

Initial fields of seabed substrates are based on shear stress and sediment type (sedimentation-, transportation- or erosion area), which were related to empirical relationships to measured sediment data across Denmark. Initial values of benthic vegetation (e.g. eelgrass) and pelagic phase were estimated by running a MIKE ECO Lab model with defined initial biomass for the entire model domain for a three-year simulation period.

In DHI (2020) further details on initial model values are given.

5 Model Calibration and Validation

After set-up of the model, calibration and validation of the model are undertaken. The model **calibration** is the process of adjusting model process settings and model constants within the literature range to obtain satisfactory agreement between observations and model results in the local modelling domain. In practice, the model set-up and the model calibration are often performed iteratively, since a good comparison between observations and model results requires a well-proportioned model domain as well as adequate model forcings.

The model **validation** is the process of comparing observations and model results qualitatively and quantitatively for a different period from the calibration period, to demonstrate the suitability of the calibrated model more generally. The qualitative comparison is typically made graphically, and the quantitative comparison is usually made using specific performance (goodness of fit) measures (DHI 2019b; Erichsen & Timmermann 2017). As such, the model validation constitutes the final documentation of the model performance.

The NBH model was run for the period 2002-2016, and the entire period is used for a combined calibration and validation effort, due to lack of enough observation data for separate calibration and validation tasks. Consequently, model performance measures are presented for this period. The model results compared with observations of the different biogeochemical parameters are given for the entire period using a WEB-tool (rbmp2021-2027.dhigroup.com, Google Chrome only).

5.1 Model Calibration Procedure

Calibration of the biogeochemical NBH model is achieved by tuning model constants to optimize model results on calibration parameters compared to measured data. The constants adjusted in the calibration procedure are numerous. They include, e.g. phytoplankton growth rates, grazing rates, mortality rates (phytoplankton and zooplankton), light attenuation constants, sedimentation rates, re-suspension rates, mineralization rates (pelagic and sediment), denitrification rates (pelagic and sediment).

The key parameters to optimize in the calibration procedure include dissolved oxygen, chlorophyll-a, light attenuation, inorganic nutrients, total nitrogen and total phosphorus. After each adjustment of calibration constants, the model is run, and time-series are compared to measured data at selected stations. The procedure is iterated until model results and measured data compare in both time and space.

5.2 Presentation of Key Model Results

During the model calibration procedure, an extensive amount of data on state variables and processes is produced. To allow for smooth and homogeneous quality assurance, few standard plots and time series are generated automatically and evaluated during the baseline and scenario execution.

Examples of modelled key validation parameters are presented as 2D fields in Figure 5-1 to Figure 5-4, and illustrate the spatial variation of the validation parameters. In the following, a brief assessment of the spatial distribution of key parameters will be given.

Average concentrations of dissolved oxygen in bottom waters during 2016 range from a minimum of 0.8 – 2.2 mg/l in the deep channels to maximum values of 9-13 mg/l found in the coastal areas (Figure 5-1).




Figure 5-1 Modelled annual average bottom water concentrations of dissolved oxygen (mg/l) for 2016.

Annual average concentrations of surface chlorophyll-a during 2016 range from 0.01 $\mu\text{g/l}$ to 23 $\mu\text{g/l}$ in the Northern Belt Sea (Figure 5-2). The highest average concentrations of chlorophyll-a in the surface during 2016 are observed in the inner Vejle Fjord, Horsens Fjord and Klaundborg Fjord, in Nærå Strand and where Odense Fjord flows into the model area.




Figure 5-2 Modelled annual average surface water concentrations of chlorophyll-a (mg/l) for 2016.

The annual average concentrations of surface total nitrogen during 2016 are highest in the inner Vejle Fjord, Horsens Fjord and Klaundborg Fjord, in Nærá Strand, Norsminde Fjord and where Odense Fjord flows into the model area, with values ranging between 1.2 – 2.6 mg/l (Figure 5-3). In the open waters of the model domain, average surface concentrations of total nitrogen range between 0.1 - 0.4 mg/l.




Figure 5-3 Modelled yearly average surface water concentrations of total nitrogen (mg/l) for 2016.

Yearly average concentrations of surface total phosphorus during 2016 are between 0.02 mg/l to 0.3 mg/l with the highest values in the inner Vejle Fjord, Horsens Fjord and Norsminde Fjord and where Odense Fjord flows into the model area (Figure 5-4).




Figure 5-4 Modelled yearly average surface water concentrations of total phosphorus (mg/l) for 2016.

5.3 Model Performance

The NBH biogeochemical model was calibrated and validated against measured data (observations) on modelled ecosystem parameters at selected stations within the model domain. Figure 5-5 shows the location of 25 stations within the model domain (stations with ID only). Of the 25 stations, 12 stations had enough measurement data in the period 2002-2016 to be included in the model calibration and validation (at least one year of weekly or bi-weekly data). The biogeochemical calibration/validation parameters include dissolved oxygen (DO), chlorophyll-a (CH), light attenuation (K_d), dissolved inorganic nitrogen (DIN), dissolved inorganic phosphorus (DIP), total nitrogen (TN) and total phosphorus (TP). Generally, the NBH model compares well to the measurements in terms of model parameters (see Figure 5-1 to Figure 5-4 and Figure 5-6 to Figure 5-12), and the overall performance measure (summarized in Table 5-1 to Table 5-3) confirms a statistically good agreement between measurements and model results.




Figure 5-5 Stations within the NBH-model domain (only stations with ID).

5.3.1 Calibration/Validation at Station VEJ0005790

In the following, we present an example on the calibration/validation from the Northern Belt Sea at station VEJ0005790 and refer to rbmp2021-2027.dhigroup.com (Google Chrome only) for more details on the Danish measurement stations. The location of station VEJ0005790 is shown in Figure 5-5.

The comparison at station VEJ0005790 shows a good agreement between the measurements and the NBH model for 76% of the parameters according to the three performance measure P-Bias, Spearman Rank Correlation and CF (see Table 5-1, Table 5-2 and Table 5-3).

In Figure 5-6, measured and modelled concentrations of dissolved oxygen (DO) at station VEJ0005790 in the surface and bottom waters (here 3 m) are shown. From the figure, it is seen that for DO the variability and seasonality of the surface and bottom waters are well represented by the model. This is in agreement with the statistical performance measures (see Table 5-1 to Table 5-3), where measured and modelled DO compare 'excellent' (P-Bias) and 'very good' (Spearman Rank Correlation and CF) at station VEJ0005790.




Figure 5-6 Comparison of measured and modelled concentrations of dissolved oxygen (mg/l) at station VEJ0005790 in the surface and bottom (3 m) waters. Dots represent measurements, and the solid line shows modelled data for the entire period.

For chlorophyll-a (CH), the dynamics in seasonality are well represented by the model (see Figure 5-7). From the statistical performance measures, annual CH compares 'excellent' and 'good' according to P-Bias and CF, respectively (Table 5-1 and Table 5-3) and 'good' based on Spearman Rank Correlation (Table 5-2).




Figure 5-7 Comparison of measured and modelled concentrations of chlorophyll-a (mg/l) at station VEJ0005790 in surface and bottom (3 m) waters. Dots represent measurements, and solid lines show modelled data for the entire period.

The measured light attenuation coefficient (K_d) presented in Figure 5-8 is well represented by the NBH model. The model tends to overestimate the winter and spring values of K_d . The statistical performance for the annual period measured by P-Bias and CF for this parameter is 'good' and based on Spearman Rank Correlation the performance for K_d is 'poor' (see Table 5-1, Table 5-2 and Table 5-3).




Figure 5-8 Comparison of measured and modelled light attenuation coefficient (K_d) at station VEJ0005790 in the surface waters. Dots represent measurements, and the solid line shows modelled data for the entire period.

For dissolved inorganic nitrogen (DIN), the structure in the seasonality is well represented by the NBH model (see Figure 5-9). From the statistical performance measures (Table 5-1, Table 5-2 and Table 5-3), annual DIN compares 'excellent' (CF), 'very good' (Spearman Rank Correlation) and 'good'.




Figure 5-9 Measured and modelled concentrations of dissolved inorganic nitrogen (DIN) at station VEJ0005790 in the surface and bottom (3 m) waters. Dots represent measurements, and solid lines show modelled data for the entire period.

When comparing measured and modelled concentrations of DIP (see Figure 5-10), we see relatively similar winter concentrations and a definite drop in spring. In general, the model tends to overestimate concentrations of DIP. The model performance for annual DIP is 'good' based on the CF measures at VEJ0005790 and 'poor' according to P-Bias and Spearman Rank Correlation (Table 5-1 to Table 5-3).




Figure 5-10 Measured and modelled concentrations of dissolved inorganic phosphorus (DIP) at station VEJ0005790 in the surface and bottom (3 m) waters. Dots represent measurements, and solid lines show modelled data for the entire period.

In Figure 5-11, comparisons of measured and modelled total nitrogen (TN) at station VEJ0005790 in surface water and bottom (3 m) water are shown. The surface and bottom water concentrations of TN are well represented by the NBH model. From the statistical performance measures (see Table 5-1 to Table 5-3), measured and modelled TN compare 'very good' (Spearman Rank Correlation and CF) and 'good' (P-Bias and CF) at station VEJ0005790.




Figure 5-11 Comparison of measured and modelled concentrations of total nitrogen (TN) at station VEJ0005790 in surface and bottom (3 m) waters. Scatter data represent measurements, and solid lines show modelled data for the entire period.

The seasonal dynamics in TP predicted by the model (see Figure 5-12) compare well with measured data on TP, with a tendency to overestimate. According to the three quality measures, the model performance for TP is 'excellent' (P-Bias), 'good' (CF) and 'poor' (Spearman Rank Correlation).




Figure 5-12 Comparison of measured and modelled concentrations of total P (TP) at station VEJ0005790 in surface and bottom (3 m) waters. Scatter data represent measurements, and solid lines show modelled data for the entire period.

5.3.2 General Calibration/Validation for Danish Waters

For the calibration/validation period (2002-2016) 12 out of the 25 stations had a sufficient amount of measurement data (at least one year of weekly or bi-weekly data) to be included in the model performance analysis (stations listed in, e.g. Table 5-1). Figure 5-5 show the different Danish locations with measurements on ecosystem parameters (chlorophyll-a (CH), light attenuation (K_d), dissolved oxygen (DO), dissolved inorganic phosphorus (DIP), dissolved inorganic nitrogen (DIN), total nitrogen (TN), and total phosphorus (TP)) during the period 2002-2016. Time series data are presented using the WEB-tool (<http://rbmp2021-2027.dhigroup.com>).

In Table 5-1 to Table 5-3, the model performance is evaluated according to DHI (2019b) based on three performance measures: P-Bias, Spearman Rank Correlation and CF. In the tables, colour codes are included to highlight the overall model performance as 'excellent', 'very good', 'good' or 'poor'. For the biogeochemical model covering the Northern Belt Sea, we aim at 'excellent', 'very good' or 'good' model performance for 75% of the data sets on measures. All model performances (both annual and summer/winter) evaluated against the three different quality measures at 12 stations were found to be 'excellent', 'very good' or 'good' for 76% of the measurements (see Figure 5-13 and Table 5-1 to Table 5-3). The annual model performance was found to be 'excellent', 'very good' or 'good' in 81% of the measurements.




Figure 5-13 Bar chart illustrating all model performances evaluated against three different quality measures and all parameters. The vertical line indicates the aim of 75% being 'excellent', 'very good' or 'good'.

According to P-Bias (Table 5-1), the model meets 'excellent', 'very good' or 'good' in 70% of all measurements (including specific winter and summer evaluations) and 75% when evaluating the annual measurements only. A 'good' model performance measured by P-Bias for summer and winter measurements indicates that the predicted absolute values of summer chlorophyll-a,

summer light attenuation, and winter inorganic nutrient concentrations correspond well to the observed values. On average, P-bias evaluates the model performance for dissolved oxygen to be 'excellent'; for TN and TP the model performance is on average 'very good', and for K_d and DIN the average performance is 'good'. The model performance for CH and DIP is on average 'poor'.

From the quality measure Spearman Rank Correlation (Table 5-2) the model performance meets 'excellent', 'very good' or 'good' in 58% of all measurements (including specific winter and summer evaluations) and 67% in the annual measurements. A good yearly correlation obtained from the Spearman Rank measure indicates a good seasonal correlation, where the predicted dynamics correspond well with the observed seasonal variability. On average, the Spearman Rank Correlation evaluates the model performance for dissolved oxygen and DIN to be 'very good'. For DIP, TN and TP, the average performance is 'good'. The average model performance for all CH and K_d evaluated from the Spearman Rank Correlation is on average 'poor'.

According to the performance measure CF (Table 5-3), the model meets 'excellent', 'very good' or 'good' in 100% of all measurements at the 12 stations. On average, the Cost Function evaluates DO, DIN, TN, TP and CH to be 'very good'; for K_d and DIP the average model performance is 'good'.

Table 5-1 Review of model performance at station P-Bias (%) based on measured and modelled data for the validation period 2002-2016. Blue colour indicates an 'excellent' model ($|x| \leq 10\%$), dark green indicates a 'very good' model ($10 < |x| < 20\%$), light green indicates a 'good' model ($20 < |x| < 40\%$), and yellow indicates a 'poor' model ($|x| > 40\%$).

Station	TN	TP	DIP		DIN		CH		DO	K _d		Number of observations	
	Annual	Annual	Annual	Winter ^a	Annual	Winter ^a	Annual	Summer ^b	Annual	Annual	Summer ^c	Annual	Summ/Wint
ARH170002	26.7	27.9	100.1	95.7	81.8	65.7	-59.4	-39.3	-2.7	-27.3	-31.6	[302-553]	[125-216]
ARH170006	27.6	19.9	74.6	73.3	68.9	56.2	-48.2	-26.3	-3.6	-18.8	-22.2	[590-1074]	[224-392]
ARH170084	6.0	1.9	39.2	41.7	51.6	48.0	-63.5	-45.6	2.7	-24.2	-28.3	[76-301]	[23-136]
ARH250032	45.3	31.5	-49.2	12.7	39.5	3.3	-90.2	-91.2	16.2	48.8	55.1	[42-92]	[17-41]
FYN0018002	23.5	-32.4	-49.0	102.1	65.3	24.3	-5.9	45.4	16.8	-34.5	-34.5	[39-62]	[10-38]
FYN0018825	-7.2	-1.5	101.7	59.4	0.2	-3.3	-62.5	-80.5	6.4	9.4	-6.6	[133-361]	[62-107]
FYN0018843	-7.2	2.5	196.5	94.7	29.3	-1.1	-89.5	-94.9	4.1	-12.9	-30.0	[112-186]	[35-75]
VEJ0004273	28.0	4.0	33.3	46.7	36.3	27.2	-14.7	26.3	2.1	-18.7	-18.0	[262-370]	[60-192]
VEJ0005790	27.9	4.1	41.2	77.9	35.2	33.6	-3.5	21.9	2.6	-23.9	-25.5	[307-864]	[181-305]
VEJ0006489	16.8	-11.1	53.6	52.1	35.7	24.1	-49.0	-36.5	-6.7	-30.2	-34.6	[113-513]	[111-186]
VSJ41007	-4.5	-2.9	11.1	29.7	-7.3	5.4	-29.2	-34.3	4.8	1.1	-1.8	[161-328]	[72-128]
VSJ41008	0.9	-32.2	-33.8	11.4	12.7	25.6	-27.6	-43.5	21.6	-4.5	-1.4	[105-220]	[46-82]

^a Jan, Feb, Dec

^b May-Sep

^c Mar-Sep

Table 5-2 Review of model performance at station Spearman Rank Correlation (no unit) based on measured and modelled data for the validation period 2002-2016. Blue colour indicates an 'excellent' model (≥ 0.90), dark green indicates a 'very good' model (0.89-0.60), light green indicates a 'good' model (0.59-0.30), and yellow indicates a 'poor' model (< 0.30).

Station	TN	TP	DIP		DIN		CH		DO	K _d		Number of observations	
	Annual	Annual	Annual	Winter ^a	Annual	Winter ^a	Annual	Summer ^b	Annual	Annual	Summer ^c	Annual	Summ/Wint
ARH170002	0.39	0.49	0.58	0.33	0.74	0.60	0.17	0.19	0.80	0.13	0.19	[302-553]	[125-216]
ARH170006	0.39	0.51	0.60	0.23	0.75	0.53	0.25	0.38	0.80	0.21	0.23	[590-1074]	[224-392]
ARH170084	0.40	0.67	0.65	0.52	0.77	0.80	0.37	0.04	0.75	0.15	0.09	[76-301]	[23-136]
ARH250032	0.59	0.28	0.23	0.17	0.76	0.49	-0.04	-0.04	0.75	0.33	0.46	[42-92]	[17-41]
FYN0018002	0.51	0.37	0.09	-0.47	0.76	0.66	0.50	0.36	0.89	0.12	0.14	[39-62]	[10-38]
FYN0018825	0.39	0.26	0.43	0.32	0.77	0.60	0.04	0.04	0.83	-0.26	-0.31	[133-361]	[62-107]
FYN0018843	0.32	0.42	0.31	0.45	0.67	0.61	-0.11	-0.05	0.87	-0.21	-0.36	[112-186]	[35-75]
VEJ0004273	0.55	0.30	0.53	0.04	0.69	0.53	0.25	0.20	0.86	0.26	0.20	[262-370]	[60-192]
VEJ0005790	0.77	0.14	0.22	0.26	0.85	0.71	0.42	0.03	0.85	0.23	0.10	[307-864]	[181-305]
VEJ0006489	0.54	0.36	0.53	0.22	0.84	0.72	0.24	0.04	0.85	0.03	-0.03	[113-513]	[111-186]
VSJ41007	0.37	0.56	0.61	0.39	0.71	0.52	0.19	-0.21	0.80	0.17	0.16	[161-328]	[72-128]
VSJ41008	0.29	0.38	0.44	0.52	0.44	0.15	0.14	-0.26	0.67	0.27	0.22	[105-220]	[46-82]

^a Jan, Feb, Dec

^b May-Sep

^c Mar-Sep

Table 5-3 Review of model performance at station Cost Function (CF, no unit) based on measured and modelled data for the validation period 2002-2016. Blue colour indicates an 'excellent' model (≤ 0.40), dark green indicates a 'very good' model (0.39-1.00), light green indicates a 'good' model (1.01-2.99), and yellow indicates a 'poor' model (≥ 3.00).

Station	TN	TP	DIP		DIN		CH		DO	K _d		Number of observations	
	Annual	Annual	Annual	Winter ^a	Annual	Winter ^a	Annual	Summer ^b	Annual	Annual	Summer ^c	Annual	Summ/Wint
ARH170002	1.13	0.99	1.20	2.22	0.87	1.64	0.52	0.79	0.41	0.96	0.97	[302-553]	[125-216]
ARH170006	1.10	0.90	1.14	2.03	0.83	1.47	0.62	0.58	0.42	1.00	1.12	[590-1074]	[224-392]
ARH170084	0.89	0.68	0.78	0.88	0.59	0.76	0.70	1.01	0.43	1.10	1.80	[76-301]	[23-136]
ARH250032	0.78	1.20	0.51	0.93	0.56	0.62	0.85	0.90	1.17	1.08	1.05	[42-92]	[17-41]
FYN0018002	0.74	0.79	0.70	2.82	0.51	0.76	0.96	1.67	0.71	0.90	0.91	[39-62]	[10-38]
FYN0018825	0.66	0.80	1.46	1.68	0.33	0.67	1.08	1.41	0.45	1.03	0.93	[133-361]	[62-107]
FYN0018843	0.77	0.88	2.54	1.93	0.43	0.71	0.70	0.92	0.47	1.25	1.33	[112-186]	[35-75]
VEJ0004273	0.84	0.68	0.92	1.80	0.45	0.80	0.75	1.02	0.36	0.92	0.93	[262-370]	[60-192]
VEJ0005790	0.54	1.02	1.30	1.80	0.36	0.65	0.90	1.11	0.44	1.05	1.13	[307-864]	[181-305]
VEJ0006489	0.63	0.60	1.15	1.37	0.48	0.64	1.00	1.23	0.47	1.11	1.40	[113-513]	[111-186]
VSJ41007	0.73	0.67	0.76	1.50	0.54	0.75	0.58	1.03	0.45	1.00	1.11	[161-328]	[72-128]
VSJ41008	0.83	0.69	0.87	1.11	0.86	1.14	0.88	1.15	0.73	0.97	1.42	[105-220]	[46-82]

^a Jan, Feb, Dec

^b May-Sep

^c Mar-Sep

6 Conclusion

This technical note shows that the model performance for the biogeochemical model covering the Northern Belt Sea meets the performance measure 'excellent', 'very good' or 'good' for 76% of the annual measures and 81% for both yearly and summer/winter measurements evaluated against three quality measures. The ambition is to meet the above criteria in 75% of all measures for all parameters and all stations (lumped). Hence, we conclude that the biogeochemical model covering the Northern Belt Sea is well suited for modelling scenarios as part of the overall development of mechanistic models towards the RBMP 2021-2027.

7 References

- DHI (2019a). MIKE 21 & MIKE 3 Flow Model FM. Hydrodynamic and Transport Module. Scientific Documentation (http://manuals.mikepoweredbydhi.help/2019/Coast_and_Sea/MIKE_321_FM_Scientific_Doc.pdf).
- DHI (2019b). Development of Mechanistic Models. Assessment of Model Performance. DHI technical report (project no. 11822245).
- DHI (2019c). Development of Mechanistic Models. Short Technical Description of the Biogeochemical Models Applied for the Mechanistic Model Development. DHI technical report (project no. 11822245).
- DHI (2019d). Development of Mechanistic Models. Mechanistic Model for the Northern Belt Sea. Hydrodynamic model documentation. DHI report (project no. 11822245).
- DHI (2019e). Udvikling af Mekanistiske Modeller. Satellitbaseret bathymetri i Danmark. DHI Gras technical report (project no. 11822245).
- DHI (2020). Development of Mechanistic Models. Short Technical Description of the Biogeochemical Models Input Data. DHI technical report (project no. 11822245).
- Erichsen AC, Møhlenberg F, Timmermann K, Christensen JPA & Göke C (2019). Gennemgang af grundlaget for afgrænsning, karakterisering og typeinddeling af kystvande i vandområdeplanerne. Rapport. <https://mst.dk/media/187927/afgraensning-karakterisering-typologi.pdf>.
- Erichsen AC (Ed.), Timmermann K (Ed.), Christensen JPA, Kaas H, Markager S, Møhlenberg F (2017). Development of models and methods to support the Danish River Basin Management Plans. Scientific documentation. Aarhus University, Department of Bioscience and DHI, 191 pp.
- Herman P, Newton A, Schernewski G, Gustafsson B & Malve O (2017). International Evaluation of the Danish Marine Models. Miljøstyrelsen (https://mfvm.dk/fileadmin/user_upload/MFVM/Nyheder/Bilag_1_Evalueringsrapport_om_de_danske_kvaelstofmodeller__10._oktober_2017-2.pdf).